

# TCCC Journal Watch

Aslani A, Ng S, Hurley M, et al: Accuracy of identification of the cricothyroid membrane in female subjects using palpation: an observational study. *Anesth Analg* 2012;Epub ahead of print

Backer D, Aldecoa C, Njimi H, Vincent J: Dopamine versus norepinephrine in the treatment of septic shock: a meta-analysis. *Crit Care Med* 2012;40:725-730

Baker M: Military medical advances resulting from the conflict in Korea, Part I: systems advances that enhanced patient survival. *Mil Med* 177;423-429

Baker M: Military medical advances resulting from the conflict in Korea, Part II: historical clinical accomplishments. *Mil Med* 2012;177:430-435

Baugh C, Stamm J, Riley D, et al: Chronic traumatic encephalopathy: neurodegeneration following repetitive concussive and subconcussive brain trauma. *Brain Imaging Behav* 2012;Epub ahead of print

Bieneck D, Charlton D: The effect of simulated field storage conditions on the accuracy of rapid user-friendly blood pathogen detection kits. *Mil Med* 2012;177:583-588

Bisanzo M, Nichols K, Hammerstedt H, et al: Nurse-administered ketamine sedation in an emergency department in rural Uganda. *Ann Emerg Med* 2012;59:268-275

Blissett P: Controversies in the management of adults with severe traumatic brain injury. *AACN Adv Crit Care* 2012;23:188-203

Bozek J, Zimmerman S: The use of tranexamic acid may reduce the risk of death in bleeding trauma patients. *Ann Emerg Med* 2012;59:223-224

Breeze J, Allanson-Bailey L, Hunt N, et al: Mortality and Morbidity from combat neck injury. *J Trauma Acute Care Surg* 2012;72:969-974

Brown K, Guthrie H, Ramasamy A, et al: Modern military surgery: lessons from Iraq and Afghanistan. *J Bone Joint Surg Br* 2012;94:536-543

Burns T, Stinner D, Mack A, et al: Microbiology and injury characteristics in severe open tibia fractures from combat. *J Trauma Acute Care Surg* 2012;72:1062-1067

Byard R, Gabrielian L, Helps S, et al: Further investigations into the speed of cerebral swelling following blunt cranial trauma. *J Forensic Sci* 2012;Epub ahead of print

Carness J, Russell J, MeLima R, et al: Fluid resuscitation using the intraosseous route: infusion with lactated ringer's and hetastarch. *Mil Med* 2012;177:222-228

Chan T, Conovaloff C, Mehran A: Hetastarch use in bariatric surgery: word of caution. *Am Surg* 2012;78:278-279

Cleiman P, Nemeth J, Vetere P: A significant cervical spine fracture: think of the airway. *J Emerg Med* 2012;42:e23-e25

D'Orazio J, Fischel J: Recurrent respiratory depression associated with fentanyl transdermal patch gel reservoir ingestion. *J Emerg Med* 2012;543-548

Doran C, Doran C, Woolley T, et al: Targeted resuscitation improves coagulation and outcome. *J Trauma Acute Care Surg* 2012;72:835-843

Effgen G, Hue C, Vogel E, et al: A multiscale approach to blast neurotrauma modeling: part II: methodology for inducing blast injury to in vitro models. *Front Neurol* 2012;3:23

Fitzgibbons P, DiGiovanni C, Hares S, Akelman E: Safe tourniquet use: a review of the evidence. *J Am Acad Orthop Surg* 2012;20:310-319

Hetzler MR: Damage control and point of injury care: extending the care continuum to military prehospital providers. *JSOM* 2012;12:1-10

Kramer C, Freeman W, Larson J, et al: Therapeutic hypothermia of severe traumatic brain injury a critically appraised topic. *Neurologist* 2012;18:173-177

Krebsbach M, Muci-Küchler K, Hinz B: Effect of initial projectile speed on contamination distribution in a lower extremity surrogate "wound track". *Mil Med* 2012;177:573-582

Kumar A, Suneja M: Hetastarch-induced osmotic nephrosis. *Anesthesiology* 2012;Epub ahead of print

Lafrance J, Miller D: Dispensed selective and nonselective nonsteroidal anti-inflammatory drugs and the risk of moderate to severe hyperkalemia: a nested case-control study. *Am J Kidney Dis* 2012;Epub ahead of print

Laiet K, Laiet J, King B, et al: Prehospital burn management in a combat zone. *Prehosp Emerg Care* 2012;16:273-276

Levi M: Should antifibrinolytics be given in all patients with trauma? Curr Opin Anaesthesiol 2012;Epub ahead of print

Li T, Lin X, Zhu Y, et al: Short-term mild hypothermia can increase the beneficial effect of permissive hypotension on uncontrolled hemorrhagic shock in rats. Anesthesiology 2012;Epub ahead of print

Mabry R: An analysis of battlefield cricothyrotomy in Iraq and Afghanistan. J Spec Oper Med 2012;12:17-23

Minel J, Cuschieri J, Sperry J, et al: The changing pattern and implications of multiple organ failure after blunt injury with hemorrhagic shock. Crit Care Med 2012;40:1129-1135

Moawad, F, Wilson R, Kunar M, Hartzell J: Role of the battalion surgeon in the Iraq and Afghanistan War. Mil Med 2012;177:412-416

Mrvos R, Feuchter A, Katz K, et al: Whole fentanyl patch ingestion: a multi-center case series. J Emerg Med 2012;42:549-552

Murad I, Abib S, Lima D, et al: The influence of hemorrhagic shock on ventilation through needle cricothyroidotomy in pigs. Am J Emerg Med 2012; Epub ahead of print

Nagao T, Kinoshita K, Sakurai A, et al: Effects of bag-mask versus advanced airway ventilation for patients undergoing prolonged cardiopulmonary resuscitation in pre-hospital setting. J Emerg Med 2012;42:162-170

Napolitano L: Guideline compliance in trauma: evidence-based protocols to improve trauma outcomes? Crit Care Med 2012;40:990-991

Neill A, Anderson P: Observational cadaveric study of emergency bystander cricothyroidotomy with a ballpoint pen by untrained junior doctors and medical students. Emerg Med J 2012;Epub ahead of print

O'Connell K, Littleton-Kearney M, Bridges E, Bibb S: Evaluation the Joint Theater Trauma Registry as a data source to benchmark casualty care. Mil Med 2012;177:564-552

O'Shea M: Acinetobacter in modern warfare. Int J Antimicrob Agents 2012;Epub ahead of print

Paix B, Griggs W: Emergency surgical cricothyroidotomy: 24 successful cases leading to a simple 'scalpel-finger-tube' method. Emerg Med Australas 2012;24:23-30

Pape A, Kutschker S, Kertscho H, et al: The choice of the intravenous fluid influences the tolerance of acute normovolemic anemia in anesthetized domestic pigs. Crit Care 2012;Epub ahead of print.

Pearson W, Ovalle F, Faul M, Sasser M: A review of traumatic brain injury trauma center visits meeting physiologic criteria from the American College of Surgeons Committee on Trauma and Center for Disease Control and Prevention Field Triage Guidelines. Prehosp Emerg Care 2012;Epub ahead of print

Rajan J, de Mello W: An alternative approach to needle thoracostomy for tension pneumothorax. Injury 2012;Epub ahead of print.

Sadaka F, Veremakis C: Therapeutic hypothermia for management of intracranial hypertension in severe traumatic brain injury: a systematic review. Brain Inj 2012;Epub ahead of print

Saulle M, Greenwald B: Chronic traumatic encephalopathy: a review. Rehabil Res Pract 2012;Epub ahead of print

Schochl H, Maegele M, Solomon C, et al: Early and individualized goal-directed therapy for trauma-induced coagulopathy. Scand J Trauma Resusc Emerg Med 2012;Epub ahead of print

Smith J, Fawcett R, Randalls B: The use of recombinant activated Factor VII in a patient with penetrating chest trauma and ongoing pulmonary hemorrhage. Mil Med 2012;177:614-616

Sobottka I, Wegscheider K, Balzer L, et al: Microbiological analysis of a prospective, randomized, double-blind trial comparing moxifloxacin and clindamycin in the treatment of odontogenic infiltrates and abscesses (MOCLI study). Antimicrob Agents Chemother 2012;Epub ahead of print

Stratton S: Rethinking out-of-hospital intravenous access. Ann Emerg Med 2012;59:304-306

Tao K, Shenbagamurthi S: Coup and contrecoup head injury resulting in anosmia. J Emerg Med 2012;42:180-181

Tuma M, Stansbury L, Stein D, et al: Induced hypothermia after cardiac arrest in trauma patients: a case series. J Trauma 2012;71:1524-1527

Waibel K, Haney B, Moore M, et al: Safety of chitosan bandages in shellfish allergic patients. Mil Med 2011;176:1153-1156

Walcott B, Nahed B, Sheth S, et al: Bilateral hemicraniectomy in non-penetrating traumatic brain injury. J Neurotrauma 2012;Epub ahead of print

Waldman M, Richman A, Shapira S: Tactical Medicine - the Israeli revised protocol. Mil Med 2012;177:52-55

Wildgruber M, Rummeny E: Bilateral tension pneumothorax. Emerg Med J 2012;Epub ahead of print

Yamagiwa T, Morita S, Inokuchi S: Re: an alternative approach to needle thoracostomy for tension pneumothorax. Injury 2012;Epub ahead of print

Yuan F, Ding J, Chen H, et al: Predicting progressive hemorrhagic injury after traumatic brain injury: derivation and validation of a risk score based on admission of characteristics. J Neurotrauma 2012;Epub ahead of print

## **February 2012**

Aboud E, Krisht A, O'keeffe T, et al: Novel simulation for training trauma surgeons. *J Trauma* 2011;71:1484-1490

Ackland H, Cameron P, Varma D, et al: Cervical spine resonance imaging in alert, neurologically intact trauma patients with persistent midline tenderness and negative computed tomography results. *Ann Emerg Med* 2011;58:521-530

Ahearn A: Perspectives "Always the Lessons". *Mil Med* 2012 177:4-5

Ali J, Sorvari A, Haskin D, et al: Potential role of the advanced surgical skills for exposure in trauma (ASSET) course in Canada. *J Trauma* 2011;71:1491-1493

Beckett A, Savage E, Pannell D, et al: Needle decompression for tension pneumothorax in Tactical Combat Casualty Care: do catheters placed in the midaxillary line kink more often than those in the midclavicular line? *J Trauma* 2011;71:S408-S412

Bragge P, Chau M, Pitt V, et al: An overview of published research about the acute care and rehabilitation of traumatic brain injured and spinal cord injured patients. *J Neurotrauma* 2011;Epub ahead of print

Brown C, Sowery L, Curry E, et al: Recombinant Factor VIIa to correct coagulopathy in patients with traumatic brain injury presenting to outlying facilities before transfer to the regional trauma center. *Am Surg* 2012;78:57-60

Bruner D, Gustafson C, Visintainer C: Ballistic injuries in the emergency department. *Emerg Med Pract* 2011;13:1-30

Capp R, Chang Y, Brown D: Effective antibiotic treatment prescribed by emergency physicians in patients admitted to the intensive care unit with severe sepsis or septic shock: where is the gap? *J Emerg Med* 2010;41:573-580

Chi J, Knudson M, Vasaar M, et al: Prehospital hypoxia affects outcome in patients with traumatic brain injury a prospective multicenter study. *J Trauma* 2011;2006:1134-1141

Childers R, Tolentino J, Leasiolagi J: Tourniquets exposed to Afghanistan combat environment have decreased efficacy and increased breakage compared to unexposed tourniquets. *Mil Med* 2011;176:1400-1403

Comstock S, Pannell D, Talbot M, et al: Spinal injuries after improvised explosive device incidents: implications for tactical combat casualty care. *J Trauma* 2011;71:S413-S417

Cross JD, Johnson AE, Wenke JC, Bosse MJ, Ficke JR: Mortality in female war veterans of Operations Enduring Freedom and Iraqi Freedom. *Clin Orthop Relat Res* 2011; E-pub 10 March 2011

Csókay A, Emelifeonwu J, Fügedi L, et al: The importance of very early decompressive craniectomy as a prevention to avoid the sudden increase of intracranial pressure in children with severe traumatic brain swelling (retrospective case series). *Childs Nerv Syst* 2011;Epub ahead of print

Eastridge B, Salinas J, Wade C, Blackbourne L: Hypotension is 100mm Hg on the battlefield. *Am J Surg* 2011;202:404-408

Faridar A, Bershad E, Emiru T, et al: Therapeutic hypothermia in stroke and traumatic brain injury. *Front Neurol* 2011;2:80

Finfer S: Hydroxyethyl starch in patients with trauma. *Br J Anaesth* 2012;108:160-161

Foerch C, You Z, Wang H, et al: Traumatic brain injury during warfarin anticoagulation- an experimental study in mice. *J Neurotrauma* 2011;Epub ahead of print

Friess S, Naim M, Kilbaugh T, et al: Premedication with meloxicam exacerbates intracranial haemorrhage in an immature swine model of non-impact inertial head injury. *Lab Anim* 2012; Epub ahead of print

Godier A, Samama C, Susen S: Plasma/platelets/red blood cell ratio in the management of the bleeding traumatized patient: does it matter? *Curr Opin Anaesthesiol*;Epub ahead of print

Graham D, Eastman A, Aldy K, et al: Outcomes and long term follow-up after emergent cricothyroidotomy: is routine conversion to tracheostomy necessary? *Am Surg* 2011;77:1707-1711

Green J, Karras D: Update on emerging infections: news from the Centers for Disease Control and Prevention. *Ann Emerg Med* 2012;59:53-54

Hamaekers A, Henderson J: Equipment and strategies for emergency tracheal access in the adult patient. *Anaesthesia* 2011;66:65-80

Harcke H, Crawley G, Mabry R, Mazuchowski E: Placement of tibial intraosseous infusion devices. *Mil Med* 2011;176:824-827

Ho A, Dion P, Yeung J, et al: Prevalence of survivor bias in observational studies on fresh frozen plasma: erythrocyte ratios in trauma requiring massive transfusion. *Anesthesiology* 2012;Epub ahead of print.

Inaba K, Branco B, Eckstein M, et al: Optimal positioning for emergent needle thoracostomy: a cadaver-based study. *J Trauma* 2011;71:1099-1103

Inaba K, Branco B, Rhee P: Impact of the duration of platelet storage in critically ill trauma patients. *J Trauma* 2011;71:1766-1774

Isayama K, Kakatani T, Tsuda M, Hirakawa A: Current status of establishing a venous line in CPA patients by emergency life-saving technicians in the prehospital setting in Japan and a proposal for intraosseous infusion. *Int J Emerg Med* 2012;Epub ahead of print

Jacobs S, Taylor M, Parker C: Changes in surgical workload at the JF Med Gp Role 3 Hospital, Camp Bastion, Afghanistan, November 2008-2010. *Injury*;Epub ahead of print

Kalwerisky K, Davies B, Mihora L, et al: Use of the Boston Ocular Surface Prosthesis in the management of severe periorbital thermal injuries: a case series of 10 patients. *Ophthalmology* 2011;Epub ahead of print

Kang D, Lehman R, Carragee E: Wartime spine injuries: understanding the improvised explosive device and biophysics of blast trauma. *Spine J* 2011;Epub ahead of print

Kauver D, Sarfati M, Kraiss L: Intraoperative blood product resuscitation and mortality in ruptured abdominal aortic aneurysm. *J Vasc Surg* 2012;Epub ahead of print

King D: Thirty consecutive uses of a hemostatic bandage at a US Army combat support hospital and forward surgical team in Operation Iraqi Freedom. *J Trauma* 2011;71:1775-1778

King D, Ogilvie M, Velmahos G, et al: Emergent cricothyroidotomies for trauma: training considerations. *Am J Emerg Med* 2011;Epub ahead of print

Kragh J, Litterel M, Jones J, et al: Battle casualty survival with emergency tourniquet use to stop limb bleeding. *J Emerg Med* 2011;41:590-597

Kragh J, O'Neill M, Walters T, et al: The military emergency tourniquet program's lessons learned with devices and designs. *Mil Med* 2011;176:1144-1152

Johnson A: Platelet transfusion: an unnecessary risk for mild traumatic brain injury patients on antiplatelet therapy. *J Emer Med* 2011;41:744-745

Lairet K, Laiet J, King B, et al: Prehospital burn management in a combat zone. *Prehosp Emerg Care* 2011;Epub ahead of print

Ley E, Singer M, Clond M, et al: Elevated admission systolic blood pressure after blunt trauma predicts delayed pneumonia and mortality. *J Trauma* 2011;71:1689-1693

Li G, Bai X, Li C, et al: In vivo antibacterial activity of chinfloxacin, a new fluoroquinolone antibiotic. *J Antimicrob Chemother* 2011;Epub ahead of print

Lundy J, Cancio L, King B, et al: Experience with the use of close-relative allograft for the management of extensive thermal injury in local national casualties during Operation Iraqi Freedom. *Am J Disaster Med* 2011;6:319-324

Mace J, Park M, Mora A, et al: Differential expression of the immunoinflammatory response in trauma patients: burn vs. non-burn. *Burns* 2011;Epub ahead of print.

Maehata Y, Esaki M, Morishita T, et al: Small bowel injury induced by selective cyclooxygenase-2 inhibitors: a prospective, double-blind, randomized clinical trial comparing celecoxib and meloxicam. *J Gastroenterol* 2011;Epub ahead of print

Malish R, Maxwell G, Witters R, et al: Perceptions of frontline providers on the appropriate qualifications for battalion level care in United States Army ground maneuver forces. *Mil Med* 2011;176:1369-1375

Martinaud C, Ausset S, Deshayes A, et al: Use of freeze-dried plasma in French intensive care unit Afghanistan. *J Trauma* 2011;71:1761-1765

Maureen M, York G, Hirshon J, et al: Trauma readiness training for military deployment: a comparison between a U.S. trauma center and an Air Force theater hospital in Blad Iraq. *Mil Med* 2011;176:769-767

Maurin O, Martinaud C, Boulesteix G, et al: Management of bleeding in a child with haemophilia in Africa with freeze-dried plasma. *Haemophilia* 2012;1:e38-e39

McGuire S, Marsh R, Sowin T, Robinson A: Aeromedical decision making and seizure risk after traumatic brain injury: longitudinal outcome. *Aviat Space Environ Med* 2012;83:140-143

Morrison JJ, Dubose JJ, Rasmussen TE, Midwinter M: Tranexamic acid decreases mortality following wartime injury: the Military Application of

Tranexamic acid in trauma and emergency resuscitation study (MATTERS); Archives of Surgery, in press

Otten M, Schwarte L, Oosterhuis J, et al: Hypercapnic coma due to spontaneous pneumothorax: case report and review of the literature. *J Emerg Med* 2012;42:e1-e6

Ozsarac M, Demican A, Sener S: Glass foreign body in soft tissue: possibility of high morbidity due to delayed migration. *J Emerg Med* 2011;41:e125-e128

Pandolf K, Francesconi R, Sawka M, et al: United States Army Research Institute of Environmental Medicine: warfighter research focusing of the past 25 years. *Adv Physiol Educ* 2011;35:353-360

Pannell D, Brisebois R, Talbot M, et al: Causes of death in Canadian Forces members deployed to Afghanistan and implications on tactical combat casualty care provision. *J Trauma* 2011;71:S401-S407

Penn-Barwell J, Murray C, Wenke J: Early antibiotics and debridement independently reduce infection in an open fracture model. *J Bone Joint Surg Br* 2012;94:107-112

Patel A, Pearce A: Progress in management of the obstructed airway. *Anaesthesia* 2011;66:93-100

Peterson E, Chesnut R: Talk and die revisited: bifrontal contusions and late deterioration. *J Trauma* 2011;71:1588-1592

Possley D, Blair J, Freedman B, et al: The effect of vehicle protection on spine injuries in military conflict. *Spine* 2011;Epub ahead of print

Possley D, Blair J, Schoenfeld A, et al: Complications associated with military spine injuries. *Spine* 2012;Epub ahead of print

Reades R, Studnek J, Vandeventer S, Garrett J: Intraosseous versus intravenous vascular access during out-of-hospital cardiac arrest: a randomized controlled trial. *Ann Emerg Med* 2011;58:509-516

Rhee P, Pirola E, Hébert-Blouin M, et al: Concomitant traumatic spinal cord and brachial plexus injuries in adult patients. *J Bone Joint Surg Am.* 2011;93:2271-2277

Riha G, Kunio N, Hamilton G, et al: Hextend and 7.5% hypertonic saline with Dextran are equivalent to lactated ringer's in a swine model of initial resuscitation of uncontrolled hemorrhagic shock. *J Trauma* 2011;71:1755-1760

Rizoli S, Scarpelini S, Callum J: Clotting factor deficiency in early trauma-associated coagulopathy. *J Trauma* 2011;71:S427-S434

Rodway G, Muza S: Fighting in thin air: operational wilderness medicine in high Asia. *Wilderness Environ Med* 2011;22:297-303

Rona R, Jones M, Fear N, et al: Frequency of mild traumatic brain injury in Iraq and Afghanistan: are we measuring incidence or prevalence? *J Head Trauma Rehabil* 2011;Epub ahead of print

Sams R, LaBrie W, Norris J, et al: IED blast postconcussive syncope and autonomic dysregulation. *Mil Med* 2012;177:48-51

Schalk R, Schweikofler U, Lotz G: Efficacy of the EZ-IO needle driver for out-of-hospital intraosseous access—a preliminary, observational, multicenter study. *Scand J Trauma Resusc Emerg Med* 2011;19:65

Schochl H, Cotton B, Inaba K, et al: FIBTEM provides early prediction of massive transfusion in trauma. *Crit Care* 2011;15:R265

Smith S, Geske J, Morgenthaler T: Risk factors associated with delayed diagnosis of acute pulmonary embolism. *J Emerg Med* 2012;42:1-6

Swan K, Swan Jr. K, Ahearn A: Tourniquets, pressure points, extremity hemorrhage. *Mil Med* 2012;177:6-7

Taecker M: Patients with traumatic brain injury. Population-based study suggests increased risk of stroke. *J Emerg of Med* 2012;42:120-121

Tan B, Chong S, Koh Z, Ong M: EZ-IO in the ED: an observational, prospective study comparing flow rates with proximal and distal tibia intraosseous access in adults. *Am J Emerg Med* 2012;Epub ahead of print

Tran L: Role of conservative management in traumatic aortic injury: comparison of long term results of conservative, surgical, and endovascular treatment. *J Emerg Med* 2012;42:122-123

Tseng W, Shih M, Su Y, et al: The association between skull bone fractures and outcomes in patients with severe traumatic brain injury. *J Trauma* 2011;71:1611-1614

Weingart S: Blunt trauma patients require a pelvic stability examination. *J Emerg Med* 2012;42:67

Wilson R, Anzueto A, Miravitles M, et al: Moxifloxacin vs amoxicillin/clavulanic acid in outpatient AECOPD: MAESTRAL results. Eur Respir J 2011;Epub ahead of print

Zierold D, Chauvire M: Hydrogen fluoride inhalation injury because of a fire suppression system. Mil Med 2012;117:108-112

## **November 2011**

BET 3: Is ketamine a viable induction agent for the trauma patient with potential brain injury. *Emerg Med J* 2011;28:1076-1077

Arnold T, Miller M, van Wessem K, et al: Base deficit from the first peripheral venous sample: a surrogate for arterial base deficit in the trauma bay. *J Trauma* 2011;Epub ahead of print

Baker A, Topolovec-Vranic J, Michalak A, et al: Controlled blast exposure during forced explosive entry training and mild traumatic brain injury. *J Trauma* 2011;71:S472-S477

Bar-Or D, Thomas G: Beneficial effects of carbamylated erythropoietin on trauma-induced brain edema: proposed molecular mechanisms of action. *Crit Care Med* 2011;39:2191

Beckett A, Savage E, Pannell D, et al: Needle decompression for tension pneumothorax in Tactical Combat Casualty Care: do catheters placed in the mid-axillary line kink more often than those in midclavicular line? *J Trauma* 2011;71:S408-S412

Bell H, Azubike E, Haouzi P: The “other” respiratory effect of opioids: suppression of spontaneous augmented (“sigh”) breaths. *J Appl Physiol* 2011;111:1296-1303

Bennett B, Cailteux-Zevallos B, Kotora J: Cricothyroidotomy bottom-up training review: battlefield lessons learned. *Mil Med* 2011;1311-1320

Béziaud T, Chen X, Shafey N, et al: Simvastatin in traumatic brain injury: Effect of brain edema mechanisms. *Crit Care Med* 2011;39:2300-2307

Blair J, Possley D, Perfield J, et al: Military penetrating spine injuries compared with blunt. *Spine J*;Epub ahead of print

Breeze J, Watson C, Horsfall I, Clasper J: Comparing the comfort and potential military performance restriction of neck collars from the body armor of six different countries. *Mil Med* 2011;176:1274-1277

Breslow J, Monroy M, Daly J, et al: Morphine, but not trauma, sensitizes to systemic acinetobacter. *J Neuroimmune Pharmacol* 2011;Epub ahead of print

Brisebois R, Tien H: Surgical experience at the Canadian-led role 3 multinational medical unit in Kandahar, Afghanistan. *J Trauma* 2011;71:S397-S400

Bouzart P, Francony G, Thomas S, et al: Reduced brain edema and functional deficits after treatment of diffuse traumatic brain injury by carbamylated erythropoietin derivative. Crit Care Med 2011;39:2099-2105

Cahill B, Stinar T: Improving the emergency whole blood program. Mil Med 2011;1287-1291

Cannon D, McCollum J: Army Medical Department lessons learned program marks 25<sup>th</sup> anniversary. Mil Med 2011;176:1212-1214

Comstock S, Pannell D, Talbot M, et al: Spinal injuries after improvised explosive device incidents: implications for Tactical Combat Casualty Care. J Trauma 2011;71:S413-S417

Culligan W, Tien H: Tranexamic acid autoinjector for prehospital care of noncompressible hemorrhage. J Trauma 2011;71:S501-S502

Darlington D, Delgado A, Kheirabadi B, et al: Effect of hemodilution on coagulation and recombinant factor VIIa efficacy in human blood in vitro. J Trauma 2011;Epub ahead of print

Davenport R, Manson J, De'Ath H, et al: Functional definition and characterization of acute traumatic coagulopathy. Crit Care Med 2011;39:2652-2658

Dean P, O'Neill D, Sterr A: Post-concussion syndrome: prevalence after mild traumatic brain injury in comparison with a sample without head injury. Brain Inj 2011;Epub ahead of print

Devlin J, Kircher S, Kozen B, et al: Comparison of Chitoflex®, CELOX™, and QuickClot® in control of hemorrhage. J Emerg Med 2011;41:237-245

Eikermann M, Grosse-Sundrup M, Zarembra S, et al: Ketamine activates breathing and abolishes the coupling between loss of consciousness and upper airway dilator muscle dysfunction. Anesthesiology 2011;Epub ahead of print

Engels P, Rezende-Neto J, Mahroos M, et al: The natural history of trauma-related coagulopathy: implications for treatment. J Trauma 2011;71:S448-S455

Feng K, Shen L, Pang X: Case-control study of risk factors for no light perception after open-globe injury. Retina 2011;31:1988-1996

Green S: Cheerio, laddie! Bidding farewell to the Glasgow Coma Scale. Ann Emerg Med 2011;58:427-430

Haut ER, Kalish BT, Cotton BA, et al: Prehospital Intravenous Fluid Administration is Associated With Higher Mortality in Trauma Patients: A National Trauma Data Bank Analysis; Ann Surg 2011;253:371-377

Herff H, Schmittinger C, Goedecke A, et al: Automated emergency ventilation devices in a simulated unprotected airway. J Emerg Med 2011;41:246-251

Holcomb J, Zarzabal L, Michalek J, et al: Increased platelet: RBC ratios are associated with improved survival after massive transfusion. J Trauma 2011;71:S318-S328

Horodyski M, DiPaola C, Conrad B, Rechtine G: Cervical collars are insufficient for immobilizing as unstable cervical spine injury. J Emerg Med 2011;41:513-519

Huang Y, Lee T, Chen W, Wang Y: Safety of the nonabsorbable dural substitute in decompressive craniectomy for severe traumatic brain injury. J Trauma 2011;Epub ahead of print

Inaba K, Barmparas G, Ibrahim D, et al: Clinical examination is highly sensitive for detecting clinically significant spinal injuries after gunshot wounds. J Trauma 2011;Epub ahead of print

Konstantinidis A, Inaba K, Dubose J, et al: The impact of nontherapeutic hypothermia on outcomes after severe traumatic brain injury. J Trauma 2011;Epub ahead of print

Lai T, Jaeger L, Jones B, et al: Continuous peripheral nerve block catheter infections in combat-related injuries: a case report of five soldiers from Operation Enduring Freedom/Operation Iraqi Freedom. Pain Med 2011;12:1676-1681

Larson B, Netta K, Aisbett B: The effect of body armor on performance, thermal stress, and exertion: a critical review. Mil Med 2011;176:1265-1273

Lehman R, Paik H, Eckel T, et al: Low lumbar burst fractures: a unique mechanism sustained in our current overseas conflicts. Spine 2011;Epub ahead of print

Marchessault J, McKay P, Hammert W: Management of upper limb amputations. J Hand Surg Am 2011;10:1718-1726

Mathieu L, Bazile F, Barthélémy R, et al: Damage control orthopaedics in the context of battlefield injuries: The use of temporary external fixation on combat trauma soldiers. Orthop Traumatol Surg Res 2011;Epub ahead of print

McBeth P, Keaney M, Ball C, et al: Aeromobile modular critical care, resuscitation , and surgical suites for operational medicine. *J Trauma* 2011;71:S494-S500

McLaughlin T, Hennecke P, Garraway N, et al: A predeployment trauma team training course creates confidence in teamwork and clinical skills: a post-Afghanistan deployment validation study of Canadian forces healthcare personnel. *J Trauma* 2011;71:S487-S493

Metterlein T, Frommer M, Ginzkey C, et al: A randomized trial comparing two cuffed emergency cricothyrotomy devices using a wire-guided and a catheter-over-needle technique. *J Emerg Med* 2011;31:326-332

Mizushima Y, Ueno M, Watanabe H, et al: Discrepancy between heart rate and makers of hypoperfusion is a predictor of mortality in trauma patients. *J Trauma* 2011;71:789-792

Moore L, Turgeon A, Émond M, et al: Definition of mortality for trauma center performance evaluation: a comparative study

Palmer A, Zhang N, Zhou Y, et al: Small-volume resuscitation from hemorrhagic shock using high-molecular-weight tense-state polymerized hemoglobin. *J Trauma* 2011;71:798-807

Panzer M, Bass C, Rafaels K, et al: Primary blast survival and injury risk assessment for repeated blast exposures. *J Trauma* 2011;Epub ahead of print

Possley D, Blair J, Schoenfeld A, et al: Complications associated with military spine injuries. *Spine J* 2011;Epub ahead of print

Riwoe D, Poncia H: Subclavian artery laceration: a serious complication of needle decompression. *Emerg Med Australas* 2011;23:651-653

Sanchez L, Straszewski S, Saghir A, et al: Anterior versus lateral needle decompression of tension pneumothorax: comparison by computed tomography chest wall measurement. *Acad Emerg Med* 2011;Epub ahead of print

Sandifer J, Jones A: Is progesterone therapy beneficial for acute traumatic brain injury? *Ann Emerg Med* 2011;58:541-542

Savage E, Forestier C, Withers N, Tien H, Pannel D: Tactical Combat Casualty Care in the Canadian Forces: lessons learned from the Afghan War. *Can J Surg* 2011;59:S118-S123

Schalk R, Schweigkofler U, Lotz G, et al: Efficacy of the EZ-IO® needle driver for out-of-hospital intraosseous access a preliminary, observational, multicenter study. *Scand J Trauma Resusc Emerg Med* 2011;19:65

Schneider E, Efron D, MacKenzie E, et al: Premorbid statin use is associated with improved survival and functional outcomes in older head-injured individuals. *J Trauma* 2011;71:815-819

Shalmovitz G, Hawthorne T: Intravenous ketamine in a dissociating dose as a temporizing measure to avoid mechanical ventilation in adult patient with severe asthma exacerbation. *J Emerg Med* 2011;41:492-494

Stuke L, Pons P, Guy J, et al: Prehospital spine immobilization for penetrating trauma-review and recommendations from the prehospital trauma life support executive committee. *J Trauma* 2011;71:763-770

Tribble D, Lloyd B, Weintrob A, et al: Antimicrobial prescribing practices following publication of guidelines for the prevention of infections associated with combat-related injuries. *J Trauma* 2011;71:S299-S306

Vandromme M, Melton S, Griffin R, et al: Intubation patterns and outcomes in patients with computed tomography-verified traumatic brain injury. *J Trauma* 2011;Epub ahead of print

Woodson J: Tactical Combat Casualty Care. Assistant Secretary of Defense for Health Affairs letter dtd 23 August 2011

Yoshida K, Okimoto N, Kishimoto M, et al: Efficacy and safety of moxifloxacin for community-acquired bacterial pneumonia on pharmacokinetic analysis. *J Infect Chemother* 2011;Epub ahead of print

Zafar S, Millham F, Chang Y, et al: Presenting blood pressure in traumatic brain injury: a bimodal distribution of death. *J Trauma* 2011;Epub ahead of print

## **August 2011**

Adler E: Defining the limits of resuscitative emergency department thoracotomy: a contemporary western trauma association perspective. *J Emerg Med* 2011;41:231-232

Alam H, Hamwi K, Duggan M, et al: Hemostatic and pharmacologic resuscitation: results of a long-term survival study in a swine polytrauma model. *J Trauma* 2011;70:636-645

Arav A, Natan D: Freeze drying (lyophilization) of red blood cells. *J Trauma* 2011;70:S61-S64

Auker C, McCarron R: US Navy experience with research on, and development of, hemoglobin-based oxygen carriers. *J Trauma* 2011;70:S40-S41

Barak A, Elhalel A, Pikkel J, et al: Incidence and severity of ocular and adnexal injuries during the Second Lebanon War among Israeli soldiers and civilians. *Graefes Arch Clin Exp Ophthalmolog* 2011; Epub ahead of print

Barbosa R, Rowell S, Diggs B, et al: Profoundly abnormal initial physiologic and biochemical data cannot be used to determine futility in massively transfused trauma patients. *J Trauma* 2011;71:S364-S369

Barbosa R, Rowell S, Sambasivan C, et al: A predictive model for mortality in massively transfused trauma patients. *J Trauma* 2011;71:S370-S374

Bernard A, Moore E, Moore F, et al: Post-injury resuscitation with human polymerized hemoglobin prolongs early survival: a post hoc analysis. *J Trauma* 2011;70:S34-S37

Bobek S, Bell R, Dierks E, Potter B: Tracheotomy in the unprotected airway. *J Oral Maxillofac Surg* 2011;Epub ahead of print

Brakenridge S, Phelan H, Henley S, et al: Early blood product and crystalloid volume resuscitation: risk association with multiple organ dysfunction after severe blunt traumatic injury. *J Trauma* 2011;71:S299-305

Blackbourne L, Baer D, Cestero R, et al: Exsanguination shock: the next frontier in prevention of battlefield mortality. *J Trauma* 2011;71:S1-3

Boet S, Borges B, Naik V, et al: Complex procedural skills are retained for a minimum for 1 yr after a single high-fidelity simulation training session. *Br J Anaesth* 2011; Epub ahead of print

Breeze J, Allanson-Bailey L, Hunt N, et al: Surface wound mapping of battlefield oculo-facial injury. *Injury* 2011;Epub ahead of print

Brown L, Aro S, Cohen M, the Trauma Outcome Group: A high fresh frozen plasma: packed red blood cell transfusion ratio decreases mortality in all massively transfused trauma patients regardless of admission international normalized ratio. *J Trauma* 2011;71:S358-S363

Bulger E: 7.5% Saline and 7.5% Saline/6% Dextran for hypovolemic shock. *J Trauma* 2011;70:S27-S29

Butler F: Fluid resuscitation in Tactical Combat Casualty Care: brief history and current status. *J Trauma* 2011;70:S11-S12

Cap A, Spinella P: Severity of head injury is associated with increased risk of coagulopathy in combat casualties. *J Trauma* 2011;71:S78-81

Chen Y, Haung W: Non-impact, blast induced mild TBI and PTSD: Concepts and caveats. *Brain Inj* 2011;25:641-650

Chiapponi C, Stocker U, Korner M, Ladumer R: Emergency percutaneous needle decompression for tension pneumoperitoneum. *BMC Gastroenterol* 2011; Epub ahead of print

Ching J, Wachtel T: A simple device to teach tube thoracostomy. *J Trauma* 2011;70:1564-1567

Christensen M, Banner C, Lefering R, et al: Quality of life after severe trauma: results from the global trauma trial with recombinant Factor VIIa. *J Trauma* 2011;70:1524-1531

Cockerham G, Rice T, Hewes E, et al: Closed-eye ocular injuries in the Iraq and Afghanistan wars. *N Engl J Med* 2011;362:2172-2173

Cohen M, West M: Acute traumatic coagulopathy: from endogenous acute coagulopathy to systemic acquired coagulopathy and back. *J Trauma* 2011;70:S47-S49

Coimbra R: 3% and 5% Hypertonic Saline. *J Trauma* 2011;70:S25-S26

Convertino V, Moulton S, Grudic G, et al: Use of advanced machine-learning techniques for noninvasive monitoring of hemorrhage. *J Trauma* 2011;71:S25-S32

Cotton B: Alternative fluids for prehospital resuscitation: “pharmacological” resuscitation fluids. *J Trauma* 2011;70:S30-S31

Cotton B, Faz G, Hatch Q, et al: Rapid thrombelastography delivers real-time results that predict transfusion within 1 hour of admission. *J Trauma* 2011;71:407-417

Darlington D, Delgado A, Kheirabadi B, et al: Effect of hemodilution on coagulation and recombinant Factor VIIa efficacy in human blood in vitro. *J Trauma* 2011; Epub ahead of print

De Castro G, MacPhee M, Driscoll I, et al: New hemostatic dressing (FAST Dressing) reduces blood loss and improves survival in a grade V liver injury model in noncoagulopathic swine. *J Trauma* 2011;70:1408-1412

Eastridge B, Hardin M, Cantrell, et al: Died of wounds on the battlefield; causation and implications for improving combat casualty care. *J Trauma* 2011;71:S4-S8

Fabian T: Perfluorocarbons. *J Trauma* 2011;70:S42-S44

Fowler M, Slater T, Garza T, et al: Relationships between early acute pain scores, autonomic nervous system function, and injury severity in wounded soldiers. *J Trauma* 2011;71:S87-S90

Gerhardt RT, Berry JA, Blackbourne LH: Analysis of life-saving interventions performed by out-of-hospital medical personnel. *J Trauma* 2011;71:S109-S113

Gifford S, Propper B, Eliason J: The ischemic threshold of the extremity. *Perspect Vasc Surg Endovasc Ther* 2011;Epub ahead of print

Gong W, Marks J, Sanati P, et al: Hypertonic saline resuscitation of hemorrhagic shock does not decrease *in vivo* neutrophil interactions with endothelium in the blood-brain microcirculation. *J Trauma* 2011;71:275-282

Guo J, Liu Y, Ma Y, et al: Evaluation of emergency tourniquets for prehospital use in China. *Chin J Traumatol* 2011;14:151-155

Gruenewald M, Meybohm P, Koerner S, et al: Dynamic and volumetric variables of fluid responsiveness fail during immediate postresuscitation period. *Crit Care Med* 2011;39:1953-1959

Gwinn D, Tintle S, Kumar A, et al: Blast-induced lower extremity fractures with arterial injury: prevalence and risk factors for amputation after initial limb-preserving treatment. *J Orthop Trauma* 2011; Epub ahead of print

Hall C: Randomized objective comparison of live tissue training versus simulators for emergency procedures. Am Surg 2011;77:561-565

Hamilton G, Van P, Differding J, et al: Lyophilized plasma with ascorbic acid decreases inflammation in hemorrhagic shock. J Trauma 2011;71:292-298

Hampton T: Traumatic brain injury: a growing problem among troops serving in today's war. JAMA 2011;306:477-479

Haukoos J, Byyny R, Erickson C, et al: Validation and refinement of a rule to predict emergency intervention in adult trauma patients. Ann Emerg Med 2011;58:164-171

Ho K, Honeybul S, Litton E: Delayed neurological recovery after decompressive craniectomy for severe non-penetrating traumatic brain injury. Crit Care Med 2011;Epub ahead of print

Holcomb J: Reconstitution: reverse engineering. J Trauma 2011;70:S65-S67

Holcomb J, Wade C, The Trauma Outcomes Group: Defining present blood component transfusion practices in trauma patients: papers from the trauma outcomes group. J Trauma 2011;71:S315-S315-S317

Inaba K: Freeze-dried plasma. J Trauma 2011;70:S57-S58

Kalin L, Sakles J: The effect of stylet choice on the success rate on intubation using the glidescope video laryngoscope in the emergency department. J Emerg Med 2011;41:209

Kheirabadi B, Arnaud F, McCarron R, et al: Development of a standard swine hemorrhage model for efficacy assessment of topical hemostatic agents. J Trauma 2011;71:S139-S146

Kotwal RS, Montgomery HR, Mabry RL, Butler FK, Cain JS, Kotwal BM, Mechler KK, Blackbourne LH, Champion HR, Holcomb JB: Eliminating Preventable Death on the Battlefield. Archives of Surgery 2011;146:1350-1358

Kragh JF, Murphy C, Dubick MA, Baer DG, Johnson J, Blackbourne LH. New tourniquet device concepts for battlefield hemorrhage control. AMEDD Journal; April-June 2011

Kragh J, Swan K, Smith D, et al: Historical review of emergency tourniquet use to stop bleeding. Am J Surg 2011;Epub ahead of print

Kuiper A, Trof R, Groenveld A: ScvO<sub>2</sub> Is not a universal indicator of fluid responsiveness. J Trauma 2011;71:518-519

Jones A, Jones A, Puskarich M, et al: Whole blood lactate kinetics in patients undergoing quantitative resuscitation for septic shock. *J Emerg Med* 2011;41:219

Letson H, Dobson G: Ultra-small intravenous bolus of 7.5% NaCl/Mg<sup>2+</sup> with adenosine and lidocaine improves early resuscitation outcome in the rat after severe hemorrhagic shock *in vivo*. *J Trauma* 2011;Epub ahead of print

Lissauer M, Chi A, Kramer M, et al: Association of 6% hetastarch resuscitation with adverse outcomes in critically ill trauma patients. *Am J Surg* 2011;Epub ahead of print

Liu K: Duration of red blood cell storage influences mortality after trauma. *J Emerg Med* 2011;40:608-609

López-Rojas R, Sánchez-Céspedes J, Docobo-Pérez F, et al: Pre-clinical studies of a new quinolone (UB-8902) against *Acinetobacter baumannii* resistant to ciprofloxacin. *Int J Antimicrob Agents* 2011;Epub ahead of print

MacDonald C, Johnson A, Cooper D, et al: Detection of blast-related traumatic brain injury in U.S. military. *N Engl J Med* 2011;364:2091-2100

Malinoski D, Ewing T, Patel M, et al: The natural history of upper extremity deep venous thromboses in critically ill surgical and trauma patients: what is the role of anticoagulation? *J Trauma* 2011;71:316-322

McSwain N, Barbeau J: Potential use of prothrombin complex concentrate in trauma resuscitation. *J Trauma* 2011;70:S53-S56

McSwain N, Champion H, Fabian T, et al: State of the art fluid resuscitation 2010: prehospital and immediate transition to the hospital. *J Trauma* 2011;70:S2-S10

Minie J: Is there a role for hemopure in the prehospital setting? *J Trauma* 2011;70:S38-S39

Moore F: The use of lactated ringer's in shock resuscitation: the good, the bad and the ugly. *J Trauma* 2011;70:S15-S16

Moore F, Goslar O, Coimbra R, et al: Blunt traumatic occult pneumothorax: Is observation safe? Results of a prospective AAST multicenter study. *J Trauma* 2011;70:1019-1025

Murray C, Wilkins K, Molter N: Infections complicating the care of combat casualties during operations Iraqi Freedom and Enduring Freedom. *J Trauma* 2011;71:S62-S73

Murray M: Review: the influence of armed conflict on the development of critical care medicine. Mil Med 2011;176:674-678

Ogilvie M, Ryan M, Proctor K: Hetastarch during initial resuscitation for trauma. J Trauma 2011;70:S19-S21

Patel N, Cogbill T, Kallies K, et al: Temporary abdominal closure: long-term outcomes. J Trauma 2011;70:769-774

Paydar S, Mousavi M, Akerdi A, et al: ScvO<sub>2</sub> is not a universal indicator of fluid responsiveness. J Trauma 2011;71:518

Perstein W, Larson M: Psychophysiology and brain imaging of cognition and affect following traumatic brain injury. Int J Psychophysiol 2011 ;Epub ahead of print

Pilate S, De Clercq S: Tension pneumothorax and life saving diaphragmatic rupture: a case report and review of the literature. World J Emerg Surg 2011; Epub ahead of print

Ran Y, Hadad E, Daher S, et al: QuikClot Combat Gauze for hemorrhage control in military trauma: January 2009 Israel Defense Force experience in the Gaza Strip - a preliminary report of 14 cases. Prehosp Disaster Med 2010;25:584-588

Rajasekhar A, Gowing R, Zarychanski R, et al: Survival of trauma patients after massive red blood cell transfusion using a high or low red blood cell to plasma transfusion ratio. Crit Care Med 2011;39:1507-1513

Rentas F: Cryopreserved red blood cells: summary. J Trauma 2011;70:S45-S46

Rhee P: Albumin. J Trauma 2011;70:S22-S24

Rizoli S: PlasmaLyte. J Trauma 2011;70:S17-S18

Rogers F, Osler T, Lee J, et al: In a mature trauma system, there is no difference in outcome (survival) between level I and level II trauma centers. J Trauma 2011;70:1354-1357

Rowell S, Barbosa R, Diggs B, et al: Effect of high product ratio massive transfusion on mortality in blunt and penetrating trauma patients. J Trauma 2011;71:S353-S357

Rowell S, Barbosa R, Diggs B, et al: Specific abbreviated injury scale values are responsible for the underestimation of mortality in penetrating trauma patients by the injury severity score. J Trauma 2011;71:S384-S388

Ryan M, Ogilvie M, Pereira B, et al: Heart rate variability is an independent predictor of morbidity and mortality in hemodynamically stable trauma patients. *J Trauma* 2011;70:1371-1380

Salazar-Vázquez B, Hightower C, Martini J, et al: Vasoactive hemoglobin solution improves survival in hemodilution followed by hemorrhagic shock. *Crit Care Med* 2011;39:1461-1466

Sambasivan C, Kunio N, Nair P, et al: High ratios of plasma and platelets to packed red blood cells do not affect mortality in non-massively transfused patients. *J Trauma* 2011;71:S329-S336

Schreiber M: The use of normal saline for resuscitation in trauma: *J Trauma* 2011;70:S13-S14

Shackford S: Prehospital fluid resuscitation of known or suspected traumatic brain injury. *J Trauma* 2011;70:S32-S33

Shaz B, Winkler A, James A, et al: Pathophysiology of early trauma-induced coagulopathy: emerging evidence for hemodilution and coagulation factor depletion. *J Trauma* 2011;70:1401-1407

Shuja F, Finkelstein R, Fukudome E, et al: Development and testing of low-volume hyperoncotic, hyperosmotic spray-dried plasma for the treatment of trauma-associated coagulopathy. *J Trauma* 2011;70:664-671

Singh S, Sukesan S, Kiran U, Makhija N: Gastrothorax or tension pneumothorax: a diagnostic dilemma. *J Emerg Trauma Shock* 2011;4:128-129

Spinella P, Wade C, Blackbourne L: The association of blood component use ratios with the survival of massively transfused trauma patients with and without severe brain injury. *J Trauma* 2011;71:S343-S352

Spoerke N, Michalek J, Schreiber M, The Trauma Outcomes Group: Crystalloid resuscitation improves survival in trauma patients receiving low ratios of fresh frozen plasma to packed red blood cells. *J Trauma* 2011;71:S380-S383

Stannard J, Volgas D, McGwin G, et al: Incisional negative pressure wound therapy after high-risk lower extremity fractures. *J Orthop Trauma* 2011; Epub ahead of print

Stapleton R, Martin T, Weiss N, et al: A phase II randomized placebo-controlled trial of omega-3 fatty acids for the treatment of acute lung injury. *Crit Care Med* 2011;39:1655-1662

Stein D, Hu P, Brenner M, et al: Brief episodes of intracranial hypertension and cerebral hypoperfusion are associated with poor functional outcome after severe traumatic brain injury. *J Trauma* 2011;71:364-374

Stevinson B: Effect of early plasma transfusion on mortality in patients with ruptured abdominal aortic aneurysm. *J Emerg Med* 2011;41:232-233

Tarmey N, Park C, Bartels O, et al: Outcomes following military traumatic cardiorespiratory arrest: a prospective observational study. *Resuscitation* 2011; Epub ahead of print

Taylor D, Vater G, Parker P: An evaluation of two tourniquet systems for the control of prehospital lower limb hemorrhage. *J Trauma* 2011;Epub ahead of print

Therien S, Nesbitt M, Duran-Stanton A, Gerhardt R: Prehospital medical documentation in the joint theater trauma registry: a retrospective study. *J Trauma* 2011;71:S103-S108

Thomson J: Impact of ABO-identical vs ABO-compatible non-identical plasma transfusion in trauma patients. *J Emerg of Med* 2011;40:607

Tourtier J, Jault P, Tazaroute K, et al: Tourniquets on the battlefield: could N-acetylcysteine be useful? *J Trauma* 2011;71:264

Villela N, Tsai A, Cabrales P, Intaglietta M: Improved resuscitation from hemorrhagic shock with ringer's lactate with increased viscosity in the hamster window chamber model. *J Trauma* 2011;71:418-424

Wade C, del Junco D, Holcomb J, The Trauma Outcomes Group: Variations between level I trauma centers in 24-hour mortality in severely injured patients requiring a massive transfusion. *J Trauma* 2011;71:S389-S393

Wade C, Salinas J, Eastridge B, et al: Admission hypo - or hyperthermia and survival after trauma in civilian and military environments. *Int J Emerg Med* 2011; Epub ahead of print

Washington C, Shuerer, Grubb R: Platelet transfusion: an unnecessary risk for mild traumatic brain injury patients on anti-platelet therapy. *J Trauma* 2011;71:358-363

Watters J, Van P, Hamilton G, et al: Advanced hemostatic dressings are not superior to gauze for care under fire scenarios. *J Trauma* 2011;70:1413-1419

Weingart S: Preoxygenation, reoxygenation, and delayed sequence intubation in the emergency department. *J Emerg Med* 2011;40:661-667

Weise C, Bartels U, Schultens A, et al: Using a laryngeal tube suction-device (LTS-D) reduces the "no flow time" in a single rescuer manikin study. *J Emerg Med* 2011;41:128-134

White J, Cannon J, Stannard A, et al: A porcine model for evaluating the management of non-compressible torso hemorrhage. *J Trauma* 2011;71:S131-138

Zuercher M, Kern K, Indik J, et al: Epinephrine improves 24-hour survival in a swine model of prolonged ventricular fibrillation demonstrating that early intraosseous is superior to delayed intravenous administration. *Anesth Analg* 2011;112:884-890

## **May 2011**

Allan P, Bloom B, Wanek S: Reversal of hemorrhagic shock-associated hepatic ischemia-reperfusion injury with N-acetylcysteine. *Mil Med* 2011;176:332-335

Bansal V, Fortlage D, Lee J, et al: A new clopidogrel (Plavix) point-of-care assay: Rapid determination of antiplatelet activity in trauma patients. *J Trauma* 2011;70:65-70

Baral A, Elhalel A, Pikkel J, et al: Incidence and severity of ocular and adnexal injuries during the Second Lebanon War among Israeli soldiers and civilians. *Graefes Arch Clin Exp Ophthalmol* 2011;Epub ahead of print

Basrai Z: Prehospital rapid sequence intubation improves functional outcome for patients with severe traumatic brain injury. *J Emerg Med* 2011;40:478-489

Bienek D, Charlton D: Accuracy of user-friendly blood typing kits tested under simulated military field conditions. *Mil Med* 2011;176:454-460

Boedeker B, Boedeker K, Bernhagen M, et al: Battle tracheal intubation using simulation: a multi center operational assessment of video laryngoscope. *Stud Health Technol Inform* 2011;163:74-76

Borgman M, Spinella P, Holcomb J, et al: The effect of FFP:RBC ration on morbidity in trauma patients based on transfusion prediction score. *Vox Sang* 2011;Epub ahead of print

Breeze J, Gibbons A, Shieff C, et al: Combat-related craniofacial and cervical injuries: a 5 year review from the British military. *J Trauma* 2011;Epub ahead of print

Breeze J, McVeigh K, Lee J, et al: Management of maxillofacial wounds sustained by British service personnel in Afghanistan. *Int J Oral Maxillofac Surg* 2011;Epub ahead of print

Brewer E, Reznikov B, Liberman R, et al: Incidence and predictors of intracranial hemorrhage after minor head trauma in patients taking anticoagulant and antiplatelet medication. *J Trauma* 2011;70:E1-E5

Buschmann C, Hunsaker J, Correns A, Tsokos M: Blunt head trauma or extensive tension pneumothorax? *Forensic Sci Med Pathol* 2011;Epub ahead of print

Cababas J, Brice J, DeMaio V, et al: Field-induce therapeutic hypothermia for neuroprotection after out-of-hospital cardiac arrest: a systematic review of the literature. *J Emerg Med* 2011;40:400-409

Callcut R, Johannigman J, Kadon K, et al: All massive transfusion criteria are not created equal: defining the predictive value of individual transfusion triggers to better determine who benefits from blood. *J Trauma* 2011;70:794-801

Charoencholvanich K, Siriwananasakul P: Tranexamic acid reduces blood loss and blood transfusion after TKA: a prospective randomized controlled trial. *Clin Orthop Relat Res* 2011;Epub ahead of print

Chatpun S, Cabrales P: Cardiac systolic function recovery after hemorrhage determines survivability during shock. *J Trauma* 2011;70:787-793

Clay J, Grayson J, Zierold: Comparative testing of new haemostatic agents in swine model of extremity arterial and venous hemorrhage. *Mil Med* 2010;175:280-284

CRASH-2 Collaborators: The importance of early treatment with tranexamic acid in bleeding trauma patients: an exploratory analysis of the CRASH-2 randomized controlled trial. *Lancet* 2011;377:1096-1101

Cross J, Ficke J, Hsu J, et al: Battlefield orthopaedic injuries cause the majority of long-term disabilities. *J Am Acad Orthop Surg* 2011;19:S1-S7

David H, Shipp J: A randomized controlled trial of ketamine/propofol versus propofol alone for emergency department procedural sedation. *Ann Emerg Med* 2011;57:435-441

Day M: Intraosseous devices for intravascular access in adult trauma. *Crit Care Nurse* 2011;31:76-90

DeLorenzo R, Pfaff J: Clinical quality management in the combat zones: the good, the bad, and the unintended consequences. *Mil Med* 2011;176:375-380

Diaz J, Norris P, Gunter O, et al: Triaging to a regional acute care surgery center: distance is critical. *J Trauma* 2011;70:116-119

Dorfman J, Burns J, Green D, et al: Decompressive laparotomy for refractory intracranial hypertension after traumatic brain injury. *Neurocrit* 2011;Epub ahead of print.

DuBose J, Barmparas G, Inaba K, et al: Isolated severe traumatic brain injuries sustained combat operations: Demographics, mortality outcomes, and lessons to be learned from contrasts to civilian counterparts. *J Trauma* 2011;70:11-18

Eardley W, Brown T, Bonner T, et al: Infection in conflict wounded. *Philos Trans R Soc Lond B Biol Sci* 2011;366:204-218

Eberle B, Schnüriger B, Inaba K, et al: Thromboembolic prophylaxis with low-molecular-weight heparin in patients with blunt solid abdominal organ injuries undergoing nonoperative management: Current practice and outcomes. *J Trauma* 2011;70:141-147

Fox C, Patel B, Clouse W: Update on wartime vascular injury. *Perspect Vasc Surg Endovasc Ther* 2011;Epub ahead of print

Garwe T, Cowan L, Neas B, et al: A propensity score analysis of prehospital factors and directness of transport of major trauma patients to a level I trauma center. *J Trauma* 2011;70:120-129

Gayat E, Bodin A, Sportiello C, et al: Performance evaluation of a noninvasive hemoglobin monitoring device. *Ann Emerg Med* 2011;57:330-333

Gazin N, Auger H, Jaulin P, et al: Efficacy and safety of the EZ-IO intraosseous device: Out-of-hospital implementation of a management algorithm for difficult vascular access. *Resuscitation* 2011;82:126-129

Giraud R, Siegenthaler N, Gayet-Ageron A, et al: ScvO<sub>2</sub> as a marker to define fluid responsiveness. *J Trauma* 2011;70:802-807

Gonzalez EA, Moore FA, Holcomb JB, et al: Fresh frozen plasma should be given earlier to patients requiring massive transfusion. *J Trauma* 2007;62:112-119

Guyette F, Suffoletto B, Castillo J, et al: Prehospital serum lactate as a predictor of outcomes in trauma patients: a retrospective observational study. *J Trauma* 2011;70:782-786

Henson N, Payan J, Terk M: Tibial subacute osteomyelitis with intraosseous abscess: an unusual complication of intraosseous infusion. *Skeletal Radiol* 2011;40:239-242

Holbrook TL, Galarneau MR, Dye JL, Quinn K, Dougherty DL: Morphine use after combat injury in Iraq and incidence of post-traumatic stress disorder. *NEJM* 2010;362:110-117

Hulse E, Thomas G: Vascular access on the 21<sup>st</sup> century military battlefield. *J R Army Med Corps* 2011;156:385-390

Inaba K, DuBose J, Barmparas G, et al: Clinical examination is insufficient to rule out thoracolumbar spine injuries. *J Trauma* 2011;70:174-179

Jones R: Trauma research: A time for reflection and dialogue. *J Trauma* 2011;70:263

Koci J, Dedek T, Havel E, Trlica J: Recombinant activated Factor VII—"Lifesaving or surgeon correcting"? *J Trauma* 2011;70:523-524

Koehler D, Shipman J, Davidson M, Guillamondegui O: Is early venous thromboembolism prophylaxis safe in trauma patients with intracranial hemorrhage. *J Trauma* 2011;70:324-329

Krugh J, Wade C, Baer D, et al: Fasciotomy rates in Operations Enduring Freedom and Iraqi Freedom: association with injury severity and tourniquet use. *J Orthop Trauma* 2011;25:134-139

Krauss W, Shah S, Shah S, Thomas S: Fentanyl in the out-of-hospital setting: variable associated with hypotension and hypoxemia. *J Emerg Med* 2011;40:182-187

Leong B, Reynolds P, Tiba M, et al: Effects of a combination hemoglobin based oxygen carrier-hypertonic saline solution on oxygen transport in the treatment of traumatic shock. *Resuscitation* 2011;Epub ahead of print

Letson H, Dobson G: Ultra-small intravenous bolus of 7.5% NaCl/Mg2+ with adenosine and lidocaine improves early resuscitation outcome in the rat after severe hemorrhagic shock *in vivo*. *J Trauma* 2011;Epub ahead of print

Lewis S, Jadish S: Total intravenous anesthesia for war surgery. *J R Army Med Corps* 2010;156:301-307

Ley E, Clond M, Srour M, et al: Emergency department crystalloid resuscitation of 1.5 L or more is associated with increased mortality in elderly and non elderly trauma patients. *J Trauma* 2011;70:398-400

Littlejohn L, Devlin J, Kircher S, et al: Comparison of Celox, ChitoFlex, WoundStat, and Combat Gauze hemostatic agents versus standard gauze in control of hemorrhage in a swine model of penetrating trauma. *Acad Emerg Med* 2011;18:340-350

Lustenberger T, Talving P, Lam L, et al: Unstable cervical spine fracture after penetrating neck injury: a rare entity in an analysis of 1,069 patients. *J Trauma* 2011;70:870-872

Magnotti L, Zarzaur B, Fischer P, et al: Improved survival after hemostatic resuscitation: Does the emperor have no clothes? *J Trauma* 2011;70:97-102

MacIntyre A, Quick J, Barnes S: Hemostatic dressings reduce tourniquet time while maintaining hemorrhage control. *Am Surg* 2011;77:152-165

Markel T, Ball C, Hayward T, Gomez G: Acute aortic rupture after a trans-spinal gunshot injury. *J Trauma* 2011;70:-522

Matijevic N, Kostousov V, Wang Y, et al: Multiple levels of degradation diminish hemostatic potential of thawed plasma. *J Trauma* 2011;70:71-80

Mayer J, Heard A, Dinsmore J, Lacquiere D: Laryngospasm in an unsedated patient during elective cannula cricothyroidotomy. *Anaesthesia* 2011;66:65-66

Michalek-Sauberer A, Granegger M, Gilly H: The efficacy of spontaneous and controlled ventilation with various cricothyrotomy devices: a quantitative in vitro assessment in a model lung. *J Trauma* 2011;Epub ahead of print

Moore E, Knudson M, Burlew C, et al: Defining the limits of resuscitative emergency department thoracotomy: a contemporary western trauma association preselective. *J Trauma* 2011;70:334-339

Mowery N, Gunter O, Collier B, et al: Practice management guidelines for management of hemothorax and occult pneumothorax. *J Trauma* 2011;70:510-518

Nachuraju P, Friedman A, Friedman J, Cabrales P: Exogenous nitric oxide prevents cardiovascular collapse during hemorrhagic shock. *Resuscitation* 2011;Epub ahead of print

Orman E, Conjeevaram H, Vuppalanchi R, et al: Clinical and histopathological features of fluoroquinolone-induced liver injury. *Clin Gastroenterol Hepatol* 2011;Epub ahead of print

Patel N, Cogbill T, Kallies K, et al: Temporary abdominal closure: long-term outcomes. *J Trauma* 2011;70:769-774

Paydar S, Salahi R, Abbasy H, et al: Who should be first of management of trauma patients: trauma surgeons or emergency medicine specialist? *J Trauma* 2011;70:264

Plurad D: Blast Injury 2011;Mil Med 2011;176:276-282

Ramasamy A, Masouros S, Newell N, et al: In-vehicle extremity injuries from improvised explosive devices: current and future foci. *Philos Trans R Soc Lond B Biol Sci* 2011;366:160-170

Remick K, Dickerson J, Nessen S, et al: Transforming US Army trauma care: an evidence-based review of the trauma literature. *US Army Med Dep J* 2010;Jul-Sep:4-21

Roquilly A, Mahe P, Seguin P, et al: Hydrocortisone therapy for patients with multiple trauma 2011. *JAMA* 2011;305:1201-1209

Salazar B, Hightower C, Martini J, et al: Vasoactive hemoglobin solution improves survival in hemodilution followed by hemorrhagic shock. *Crit Care Med* 2011;Epub ahead of print

Salehpour F, Bazzazi A, Porhomayon J, Nader N: Correlation between coagulopathy and outcome in severe head trauma in neurointensive care and trauma units. *J Crit Care* 2011;Epub ahead of print

Salinas N, Brennan J, Gibbons M: Massive facial trauma following improvised explosive device blasts in Operation Iraqi Freedom. *Otolaryngol Head Neck Surg* 2011;Epub ahead of print.

Sinha R, Roxby D: Transfusion practices in massive haemorrhage in pre-intensive care. *Vox Sang* 2011;Epub ahead of print

Teixeira P, Inaba K, Barmparas G, et al: Blunt thoracic aortic injuries: an autopsy study. *J Trauma* 2011;70:197-202

Truitt M, Johnson V, Rivera M, et al: Civilian and military trauma: does civilian training prepare surgeons for the battlefield? *Am Surg* 2011;77:19-21

Tsang K, Whitfield P: Traumatic brain injury: review of current management strategies. *Br J Oral Maxillofac Surg* 2011;Epub ahead of print

Vallet B: Intravascular volume expansion: which surrogate markers could help the clinician to assess the improved tissue perfusion? *Anesth Anal* 2011;112:258-259

Vandromme M, Griffin R, Kerby J, et al: Identifying risk for massive transfusion in the relatively normotensive patient: utility of the prehospital shock index. *J Trauma* 2011;70:384-390

Wafaisade A, Lefering R, Bouillon B: Epidemiology and risk factors of sepsis after multiple trauma: an analysis 29,829 patients from the Trauma Registry of the German Society for Trauma Surgery. *Crit Care Med* 2011;39:621-628

Wafaisade A, Maegele M, Lefering R, et al: High plasma to red blood cell ratios are associated with lower mortality rates in patients receiving multiple transfusion ( $4 \leq$  red blood cell units  $< 10$ ) during acute trauma resuscitation. *J Trauma* 2011;70:81-89

Wang H, Mann N, Mears G, et al: Out-of-hospital airway management in the United States. *Resuscitation* 2011;Epub ahead of print

White J, Stannard A, Burkhardt G, et al: The epidemiology of vascular injury in the wars in Iraq and Afghanistan. *Ann Surg* 2011;Epub ahead of print

Wigginton J, Roppolo L, Pepe P: Advances in resuscitative trauma care. *Minerva Anestesiologica* 2011;Epub ahead of print

Xue F, Liu H, Xiong J, et al: Facilitating smooth insertion of an endotracheal tube into the trachea under Glidescope® video laryngoscope. *J Emerg Med* 2011;40:443-447

Zahoor A, Ubiera J, Davis G, et al: Lessons learned from casualty statistics in health care system development: Afghanistan 2008-2009. *Mil Med* 2011;176:94-98

## February 2011

Alexandrou N, Yeh B, Barbara P, et al: An innovative approach to orotracheal intubations: the Alexandrou angle of intubation position. *J Emerg Med* 2011;40:7-13

Allocock E, Spencer E, Frazer R, et al: Continuous transverses abdominis plane (TAP) block catheters in a combat surgical environment. *Pain Med* 2010;11:1426-1429

Apostolakis E, Apostolaki G, Apostolaki M, Chorti M: The reported thoracic injuries in Homer's Iliad. *J Cardiothorac Surg* 2010;5:114

Bala M, Shussman N, Avraham I, et al: The pattern of thoracic trauma after suicide terrorist bombing attacks. *J Trauma* 2010;69:1022-1029

Balzli C, Caballero A, Tang A, et al: Penetration and effectiveness of prophylactic fluoroquinolones in experimental methicillin-sensitive or methicillin-resistant *Staphylococcus aureus* anterior chamber infections. *J Cataract Refract Surg* 2010;36:2160-2167

Bernard S, Nguyen V, Cameron P, et al: Prehospital rapid sequence intubation improves functional outcome for patients with severe traumatic brain injury; a randomized controlled trial. *Ann Surg* 2010;252:959-965

Blaivas M: Point-of-care ultrasonographic deep venous thrombosis evaluation after ten minutes' training: is this offer too good to be true? *Ann Emerg Med* 2010;56:611-613

Breeze J, Horsfall I, Hepper A, Clasper J: Face, neck and eye protection: adapting body armor to counter the changing patterns of injuries on the battlefield. *Br J Oral Maxillofac Surg* 2010:Epub ahead of print

Brown J, Stassen N, Bankey P, et al: Helicopters and the civilian trauma system: national utilization patterns demonstrate improved outcomes after traumatic injury. *J Trauma* 2010;69:1030-1036

Bulger E, May W, Kerby J, et al: Out-of-hospital hypertonic resuscitation after traumatic hypovolemic shock: a randomized, placebo controlled trial. *Ann Surg* 2010;Epub ahead of print

Burns J, Baer L, Hagerman E, et al: Development and resuscitation of a sedated, mature male miniature swine severe hemorrhage model. *J Trauma* 2010;Epub ahead of print

Clifton G, Valadka A, Zygund D, et al: Very early hypothermia induction in patients with severe brain injury (the National Acute Brain Injury Study: hypothermia II): a randomized trial. Lancet Neurol 2010;Epub ahead of print

Combes X, Jabre P, Margenet A, et al: Unanticipated difficult airway management in the prehospital emergency setting: prospective validation of an algorithm. Anesthesiology 2011;114:105-110

Davies G, Lockey S: Thirteen survivors of prehospital thoracotomy for penetrating trauma: a prehospital physician-performed resuscitation procedure that can yield good results. J Trauma 2010:Epub ahead of print

Dobscha S, Campbell R, Morasco B, et al: Pain in patients with polytrauma; a systemic review. Department of Veterans Affairs 2008;VA Evidence-based synthesis program reports.

Donaldson R, Hung Y, Shanovich P, et all: Injury burden during an insurgency: the untold trauma of a infrastructure breakdown in Baghdad, Iraq. J Trauma;69:1379-1385

Eastridge B, Butler F, Wade C, et al: Field triage score (FTS) in battlefield casualties: validation of a novel triage technique in a combat environment. Am J Surg 2010;200:724-727

Granville-Chapman J, Jacobs N, Midwinter M: Pre-hospital haemostatic dressings: A systematic review. Injury 2010;Epub ahead of print.

Helling T, Davit F, Edwards K: First echelon hospital care before trauma center transfer in a rural trauma system: Does it affect outcome? J Trauma 2010;69:1362-1366

Kim K, Simon L: Transport mechanisms in oral transmucosal drug delivery: implications for pain management. Math Biosci 2010:Epub ahead of print

Kirves H, Handolin L, Niemela M, et al: Paramedics' and pre-hospital physicians' assessments of anatomic injury in trauma patients: a cohort study. Scand J Trauma Resusc Emerg Med 2010;18:60

Lustenberger T, Talving P, Kobayashi L, et al: Early coagulopathy after isolated severe traumatic brain injury: relationship with hypoperfusion challenged. J Trauma 2010;69:1410-1414

Messa M, Kupas D, Dunham D: Comparison of bougie-assisted intubation with traditional endotracheal intubation in a simulated difficult airway. Prehosp Emerg Care;15:30-33

Moon J, Chun B: Fentanyl intoxication caused by abuse of transdermal fentanyl. *J Emerg Med* 2011;40:37-40

Murphy C, Rooney S, Maharaj C, et al: Comparison of three cuffed emergency percutaneous cricothyroidotomy devices to conventional surgical cricothyroidotomy in a porcine model. *Br J Anaesth* 2011;106:57-64

Palmieri T, Sen S, Falwell K, Greenhalgh D: Blood product transfusion: Blood product transfusion: does location make a difference? *J Burn Care Res* 2010;Epub ahead of print

Paul J, Webb T, Aprahamian C, Weigelt J: Intra-abdominal vascular injuries: are we getting any better? *J Trauma* 2010;69:1393-1397

Phillips L, Proehl J, Brown L, et al: Recommendations for the use of intraosseous vascular access for emergent and non-emergent situations in various health care settings: consensus paper. *J Infus Nurs* 2010;33:346-351

Schwartz R, McManus J, Croushorn J, et al: Tactical Medicine-competency based guidelines. *Prehosp Emerg Care* 2011;15:67-82

Sethuraman K, Duong D, Mehta S, et al: Complications of tube thoracostomy placement in the emergency department. *J Emerg Med* 2011;40:14-20

Shah K, Kwong B, Hazan A, et al: Success of the gum elastic bougie as a rescue airway in the emergency department. *J Emerg Med* 2011;40:1-6

Sondeen J, Prince M, Kheirabadi B, et al: Initial resuscitation with plasma and other blood components reduced bleeding compared to hetastarch in anesthetized swine with uncontrolled splenic hemorrhage. *Transfusion* 2010;Epub ahead of print

Tomimaru Y, Wada H, Marubashi S, et al: Fresh frozen plasma transfusion does not affect outcomes following hepatic resection for hepatocellular carcinoma. *World J Gastroenterol* 2010;16:5603-5610

Van Eerd M, Mario Ouwendijk J, de Peuter M: Cost-effectiveness study comparing pharmaceutically licensed plasma for transfusion (OctaplasLG®) versus fresh frozen plasma (FFP) in critically ill patients in the UK. *Transfus Apher Sci* 2010;43:251-259

Vilensky D, MacDonald R: Communication errors in dispatch of air medical transport. *Prehosp Emerg Care* 2011;15:39-43

Weinberg J, McGwin G, Vandromme M, et al: Duration of red cell storage influences mortality after trauma. *J Trauma*;69:1427-1432

## **November 2010**

Al-Hilli A, Salih D: Early or delayed surgical treatment in compound limb fractures due to high velocity missile injuries: a 5-year retrospective study from medical city in Baghdad. *Iowa Orthop J* 2010;30:94-98

Alam H, Bice L, Butt M, et al: Testing of blood products in a polytrauma model: Results of a multi-institutional randomized preclinical trial. *J Trauma* 2009;67:856-864

Aledort L: Off label use of recombinant activated Factor VII- safe or not safe? *NEJM* 2010;363:1853-1854

Allen P, Salyer S, Dubick M, et al: Preventing hypothermia: comparison of current devices used by the U.S. Army in an in vitro warmed fluid model. *J Trauma* 2010;69 Suppl:S154-161

Aoi Y, Inagawa G, Hashimoto, et al: Airway scope laryngoscopy under manual inline stabilization and cervical collar immobilization: A crossover in vivo cinefluoroscopic study. *J Trauma* 2010;Epub ahead of print

Arabi Y, Haddad S, Tamim H, et al: Mortality reduction after implementing a clinical practice guidelines-based management protocol for severe traumatic brain injury. *J Crit Care* 2010;25:190-195

Baba M, Fujimoto J, Mizutani K, et al: Tracheal intubation using the Airway Scope (®) in two patients with difficult airway during cardiopulmonary resuscitation. *J Anesth* 2010;24:18-20

Ball C, Wyrzykowski A, Kirkpatrick A, et al: Thoracic needle decompression for tension pneumothorax: clinical correlation with catheter length. *Can J Surg* 2010;53:184-188

Bassi M, Zuercher M, Erne J, Ummenhofer W: Endotracheal tube intracuff pressure during helicopter transport. *Ann Emerg Med* 2010;56:89-93

Béchir M, Puhan M, Neff S, et al: Early fluid resuscitation with hyperoncotic hydroxyethyl starch 200/0.5 (10%) in severe burn injury. *Crit Care* 2010;14:R123

Beiser D, Wang H, Li J, et al: Plasma and myocardial visfatin expression changes are associated with therapeutic hypothermia protection during murine hemorrhagic shock/resuscitation. *Resuscitation* 2010;Epub ahead of print

Belmont P, Thomas D, Goodman G, et al: Combat musculoskeletal wounds in a US Army brigade combat team during Operation Iraqi Freedom. *J Trauma* 2010;Epub ahead of print.

Ben-Galim P, Drengel N, Mattox K, et al: Extrication collars can result in abnormal separation between vertebrae in the presence of a dissociative injury. *J Trauma* 2010;Epub ahead of print

Berger E: Nothing gold can stay? *Annals Emerg Med* 2010;56:17A-19A

Boldt, Joachim: Modern rapidly degradable hydroxyethyl starches: current concepts. *Anesth Analg* 2009;108:1574-1582

Bridges E, Biever K: Advancing critical care: Joint Combat Casualty Team and Joint Theater Trauma System. *AACN Adv Crit Care* 2010;21:260-276

Brown K, Murray C, Clasper J: Infectious complications of combat-related mangled extremity injuries in the British military. *J Trauma* 2010;69 Suppl:109-115

Butler F: Tactical Combat Casualty Care: Update 2009. *J Trauma* 2010;69 Suppl:S10-13

Byers R: An upshot of war-damage control resuscitation. *Int Emerg Nurs* 2010;Epub ahead of print

Calderbank P, Woolley T, Mercer S, et al: Doctor on board? What is the optimal skill-mix in military pre-hospital care? *Emerg Med* 2010;Epub ahead of print

Cha Y, Cho T, Suh J: Traumatic cervical cord transection without facet dislocations- a proposal of combined hyperflexion-hyperextension mechanism: a case report. *J Korean Med Sci* 2010;25:1247-1250

Chatpun S, Cabrales P: Cardiac systolic function recovery after hemorrhage determines survivability during shock. *J Trauma* 2010;Epub ahead of print

Crandall M: "Live Tissue" is not the answer. *J Trauma* 2010;69(4):998-999

Davis D, Peay J, Sise M, et al: Prehospital airway and ventilation management: a trauma score and injury severity score-based analysis. *J Trauma* 2010;69:294-301

Davis K, Rozycki G: Acute care surgery in evolution. *Crit Care Med* 2010;38:S405-S410

Doran C, Wooley T, Midwinter M: Feasibility of using rotational thromboelastometry to assess coagulation status of combat casualties in a deployed setting. *J Trauma* 2010;69 Suppl:40-48

Duchesne J, McSwain N, Cotton B, et al: Damage control resuscitation: the new face of damage control. *J Trauma* 2010;69:976-990

Elliot D, Baker P, Scott M, et al: Accuracy of surface landmark identification for cannula cricothyroidotomy. *Anaesthesia* 2010;Epub ahead of print

Fenwick R: Intraosseous approach to vascular access in adult resuscitation. *Emerg Nurse* 2010;18:22-25

Fox C, Perkins J, Kragh J, et al: Popliteal artery repair in massively transfused military trauma casualties: a pursuit to save life and limb. *J Trauma* 2010;69 Suppl:123-134

Gataa I, Muassa Q: Patterns of maxillofacial injuries caused by terrorist attacks in Iraq: retrospective study. *Int J Oral Maxillofac Surg* 2010;Epub ahead of print

Giraud R, Siegenthaler N, Gayet-Ageron A, et al: ScvO<sub>2</sub> as a marker to define fluid responsiveness. *J Trauma* 2010;Epub ahead of print

Göskel O, Aydin O, Misirligil Z, et al: Safety of meloxicam in patients with aspirin/non-steroidal anti-inflammatory drug-induced urticaria and angiodema. *J Dermatol* 2010;37(11):973-979

Govaert G, Van Helden S: Ty-raps in trauma: a novel closing technique of extremity fasciotomy wounds. *J Trauma* 2010;69:972-975

Hall A, Evans K, Pribyl S: Cold injury in the United States military population: current trends and comparison with past conflicts. *J Surg Educ* 2010;67:61-65

Hirshberg A, Frykberg E, Mattox K, Stein M: Triage and trauma workload in mass casualty: a computer model. *J Trauma* 2010;Epub ahead of print

Henning D, Smith J, Patch D, Lambert A: A comparison of civilian (National Confidential Enquiry into Patient Outcome and Death) trauma standards with current practice in a deployed field hospital in Afghanistan. *Emerg Med J* 2010;Epub ahead of print

Hubble M, Brown L, Wilfong D et al: A meta-analysis of prehospital airway control techniques part II: Alternative airway devices and cricothyrotomy success rates. *Prehosp Emerg Care* 2010;14:515-530

Kellum J, James M: Why do intensivists still use hydroxyethyl starch? Crit Care Med 2010;38:2260-2262

Kheirabadi B, Mace J, Terrazas I, et al: Clot-Inducing minerals versus plasma protein dressing for topical treatment of external bleeding in the presence of coagulopathy. J Trauma 2010;69:1062-1073

King D, Namias N, Andrews D: Coagulation abnormalities following thermal injury. Blood Coagul Fibrinolysis 2010;Epub ahead of print

Kummoona R: Management of missile injuries of the facial skeleton: primary, intermediate, and secondary phases. J Craniofac Surg 2010;21:976-981

Letson H, Dobson G: Small volume 7.5% NaCl with 6% Dextran-70 or 6% and 10% hetastarch are associated with arrhythmias and death after 60 minutes of severe hemorrhagic shock in rat in vivo. J Trauma 2010;Epub ahead of print

Levi M, Levy J, Andersen H, Truloff D: Safety of recombinant activated Factor VII in randomized clinical trials. NEJM 2010;363:1791-1800

Li L, Tomofeev I, Czosnyka M, Hutchinson P: The surgical approach to the management of increased intracranial pressure after the traumatic brain injury. Anesth Analg 2010;Epub ahead of print

Lobo D, Stanga Z, Aloysius M, et al: Effect of volume loading with 1 liter intravenous infusion of 0.9% saline, 4% succinylated gelatin (Gelofusine) and 6% hydroxyethyl starch (Voluven) on blood volume and endocrine responses: a randomized, three-way crossover study in healthy volunteers. Crit Care Med 2010;38(2):464-470

Lobo S, Stanga Z, Simpson J, et al: Dilution and redistribution effects of rapid 2-litre infusions of 0.9% (w/v) saline and 5% (w/v) dextrose on haematological parameters and serum biochemistry in normal subjects: a double blind cross-over study. Clin Sci (lond) 2010;101(2):173-179

Mabry RL, Edens JW, Pearse L, Kelly JF, Harcke H: Fatal airway injuries during Operation Enduring Freedom and Operation Iraqi Freedom. Prehospital Emergency Care 2010;14:272-277

Macgregor A, Dougherty A, Galarneau M: Injury-specific correlates of combat-related traumatic brain injury in Operation Iraqi Freedom. J Head Trauma 2010;Epub ahead of print

Madill J: In-flight thoracic ultrasound detection of pneumothorax in combat. J Emerg Med 2010;39:194-197

Marsh R, Levine A, Noble V, et al: Blunt Cardiac Rupture. *J Emerg Med* 2010;39(3):337-340

Marshall W: Resuscitation of combat casualties: unique challenges and lessons learned. *AACN Adv Crit Care* 2010;21:279-287

McLean A, Richard M, Crandall C, Marinaro J: Ultrasound determination of chest wall thickness: implications for needle thoracostomy. *Am J Emerg Med* 2010;Epub ahead of print.

McNeill M: 7,000 miles and 7 days from the battlefield. *AACN Adv Crit Care* 2010;21:307-315

Morrison J, Clasper J, Gibb I, Midwinter M: Management of penetrating abdominal trauma in the conflict environment: the role of computed tomography scanning. *World J Surg* 2010;Epub ahead of print.

Murphy G, Greenburg S: The new- generation hydroxyethyl starch solutions: the holy grail of fluid therapy or just another starch? *J Cardiovasc Anesth* 2010;24:398-393

Ogilvie M, Perira B, McKenney M, et al: First report on safety and efficacy of hetastarch solution for initial fluid resuscitation at a level 1 trauma center. *J Am Coll Surg* 2010;210:870-880.

Pehböck D, Wenzel V, Voelkel W, et al: Effects of preoxygenation on desaturation time during hemorrhagic shock in pigs. *Anesthesiology* 2010;Epub ahead of print

Pereira B, Ryan M, Ogilvie M, et al: Predeployment mass casualty and clinical trauma training for US Army forward surgical teams. *J Craniofac Surg* 2010;21:982-986

Pizzola J: MEDEVAC miscategorization. *Mil Med* 2010;175:655-658

Ramasamy A, Midwinter M, Mahoney P, Clasper J: Learning the lessons from conflict: Pre-hospital cervical spine stabilization following ballistic neck trauma. *Injury Int J Care Injured* 2010;40:1342-1345

Rangel-Castilla L, Rivera Lara L, Gopinath S, et al: Cerebral hemodynamic effects of acute hyperoxia and hyperventilation after severe traumatic brain injury. *J Neurotrauma* 2010;Epub ahead of print

Rehman S, Slemenda C, Kestner C, Joglekar S: Management of gunshot pelvic fractures with bowel injury: Is fracture debridement necessary? *J Trauma* 2010;Epub ahead of print

Roppolo L, Wigginton J, Pepe P: Intravenous fluid resuscitation for the trauma patient. *Curr Opin Crit Care* 2010;16:283-238

Sheppard F, Keiser P, Craft D, et al: The majority of US combat casualty soft-tissue wounds are not infected or colonized upon arrival or during treatment at a continental US military medical facility. *Am J Surg* 2010;200:489-495

Simmons J, White C, Eastridge B: Impact of policy change on US Army combat transfusion practices. *J Trauma* 2010;69 Suppl:S75-80

Spoerke N, Van P, Differding J, et al: Red blood cells accelerate the onset of clot formation in polytrauma and hemorrhagic shock. *J Trauma* 2010;69:1054-1061

Spoerke N, Zink K, Cho D, et al: Lyophilized plasma for resuscitation in a swine model of severe injury. *Arch Surg* 2009;144(9):829-834

Voelckel W, Convertino V, Lurie K, et al: Vasopressin for hemorrhagic shock management: revisiting the potential value in civilian and combat casualty care. *J Trauma* 2010;69 Suppl 1:69-74

Waibel B, Rotondo M: Damage control in trauma and abdominal sepsis. *Crit Care Med* 2010;38:S421-S430

Wade C, Eastridge B, Jones J, et al: Use of recombinant factor VIIa in US military casualties for a five-year period. *J Trauma* 2010;69:353-359

Westphal M, James M, Kozek-Langenecker S, et al: Hydroxyethyl Starches. *Anesthesiology* 2010;111:187-202

Willis C, Stoelwinder J, Lecky F, et al: Applying composite performance measures to trauma care. *J Trauma* 2010;69:256-262

Yoshihiro A, Gaku I, Hashimoto K, et al: Airway scope laryngoscopy under manual inline stabilization and cervical collar immobilization: a crossover in vivo cinefluoroscopic study. *J Trauma* 2010;Epub ahead of print.

## August 2010

Baer D, Dubick M, Wenke J, et al: Combat casualty care research at the U.S. Army Institute of Surgical Research. *J R Army Med Corps* 2009;155(4):327-332

Balbino M, Capone N, Prist R, et al: Fluid resuscitation with isotonic or hypertonic saline solution avoids intraneuronal calcium influx after traumatic brain injury associated with hemorrhagic shock. *J Trauma* 2010;68(4):859-864

Ball C, Wyrzykowski A, Kirkpatrick A, et al: Thoracic needle decompression for tension pneumothorax: clinical correlation with catheter length. *Can J Surg* 2010;53(3):184-188

Balzli C, Caballero A, O'Callaghan R: Comparing the effectiveness of moxifloxacin and povidone-iodine with povidone-iodine alone. *J Cataract Refract Surg* 2010;36(5):871-872

Bechir M, Puhan M, Neff S, et al: Early fluid resuscitation with hyperoncotic hydroxyethyl starch. *Crit Care* 2010; Epub ahead of print

Bourdeauz C, Brown J: Sodium bicarbonate lowers intracranial pressure after traumatic brain injury. *Neurocrit Care* 2010; Epub ahead of print

Brackney C, Diaz L, Milbrandt E, et al: Is albumin use SAFE in patients with traumatic brain injury? *Crit Care* 2010;14(2):307

Burd A: Fluid resuscitation in burns. *Burns* 2010; Epub ahead of print

Butler FK: TCCC Update 2009. *J Trauma* 2010;69:S10-S13

Cai B, Dong W, Sharpe S, et al: Survival and inflammatory responses in experimental models of hemorrhage. *J Surg Res* 2009; Epub ahead of print

Champion H, Holcomb J, Lawnick M, et al: Improved characterization of combat injury. *J Trauma* 2010;68(5):1139-1150

Chieregato A, Martino C, Pransani V, et al: Classification of traumatic brain injury: the Glasgow Coma scale is not enough. *Acta Anaesthesiol Scand* 2010; Epub ahead of print

Cho Y, Shi R, Borgens R: Chitosan produces potent neuroprotection and physiological recovery following traumatic spinal cord injury. *J Exp Biol* 2010;213(9):1513-1520

Clay J, Grayson K, Zieroid D: Comparative testing of new hemostatic agents in a swine model of extremity arterial and venous hemorrhage. Mil Med 2010;175(4):280-284

D'Alleyrand J, Dutton R, Pollak A: Extrapolation of battlefield resuscitative care to the civilian setting. J Surg Orthop Adv 2010;19(1):62-69

Dar D, Soustiel J, Zaaroor M, et al: Moderate Ringer's lactate solution resuscitation yields best neurological outcome in controlled hemorrhagic shock combined with brain injury in rats. Shock 2010;34(1):75-82

Dayan L, Zinnmann C, Stahl S, et al: Complications associated with prolonged tourniquet application on the battlefield. Mil Med 2008;173:63-66

Dewalt J: The ABCs of TBI evidence-based guidelines for adult traumatic brain injury care. JEMS 2010;35(4):54-63

Farcy DA, Peterson PL, Rabinowitz D, Scalea T: Controversies in fluid resuscitation. Emerg Med Reports 2010;14: 21 June 2010

Frith D, Brohi K: The acute coagulopathy of trauma shock: clinical relevance. Surgeon 2010;8(3):159-163

Goldberg M: Death and injury rates of U.S. Military personnel in Iraq. Mil Med 2010;175:220-226

Grape S, Schug SA, Lauer S, Schug BS: Formulations of fentanyl for the management of pain. Drugs 2010;70:57-72

Greenhalgh D: Fluid resuscitation in burns. Burns 2010; Epub ahead of print

Hadley J, Mösges R, Desrosiers M, et al: Moxifloxacin five-day therapy versus placebo in acute bacterial rhinosinusitis. Laryngoscope 2010;120(5):1057-1062

Haugk M, Stratil P, Sterz F, et al: Temperature monitored on the cuff surface of an endotracheal tube reflects body temperature. Crit Care Med 2010;38(7):1569-

Hicks R, Fertig S, Desrocher R, et al: Neurological effects of blast injury. J Trauma 2010;68(5):1257-1263

Ho K, Burrell M, Rao S; Extracranial injuries are important in determining mortality of neurotrauma. Crit Care Med 2010;38(7):1562-1586

Hospenthal DR, Crouch HK, English JF, et al: Response to infection control challenges in the deployed setting: Operations Iraqi and Enduring Freedom. *J Trauma* 2010;69:S94-S101

Kiekas P, Diamano A, Baltoplulos G, et al: Concerns about the association between temperature and mortality in trauma patients. *J Trauma* 2010;68(4):1016-1017

Kozen B, Kircher S, Henao J, et al: An alternative hemostatic dressing: comparison of Celox to Hemcon and QuickClot. *Acad Emerg Med* 2008;15:74-81

Ley E, Berry C, Mirocha J, Salim A; Mortality is reduced for heart rate 80 to 89 after traumatic brain injury. *J Surg Res* 2010;Epub ahead of print

Mercadante S, Radburch L, Davies A, et al: A comparison of intranasal fentanyl spray with oral transmucosal fentanyl citrate for the treatment of breakthrough cancer pain: an open-label, randomized, crossover trial. *Curr Med Res Opin* 2009;25(11):2805-2815

Mercer S, Guha A: Assessing the implementation of guidelines for the management of the potentially injured cervical spine in unconscious trauma patients in England. *J Trauma* 2010;68(6):1145-1450

Molin R, Hallas P, Brabrand M, Schmidt T: Current use of intraosseous infusion in Danish emergency departments: a cross-sectional study. *Scand J Trauma Resusc Emerg Med* 2010;18(1):37

Namdar T, Siemers F, Stollwerck P, et al: Increased mortality in hypernatremic burned patients. *Gen Med Sci* 2010;8:Doc 11

Nunez T, Young P, Holcomb J, et al: Creation, implementation, and maturation of a massive transfusion protocol for the exsanguinating trauma patient. *J Trauma* 2010;68(6):1498-1505

Owens D, Greenwood B, Galley A, et al: Airflow efficacy of ballpoint pen tubes: a consideration for use in bystander cricothyrotomy. *Emerg Med J* 2010;27(4):317-320

Parker P: Emergency tourniquet use. *Ann Surg* 2009;250(3):496-497

Pranger A, Alffenaar J, Wessels A, et al: Determination of moxifloxacin in human plasma, plasma ultrafiltrate, and cerebrospinal fluid by a rapid and simple liquid chromatography-tandem mass spectrometry method. *J Anal Toxicol* 2010;34(3):135-141

Propper B, Gifford S, Calhoon J, et al: Wartime thoracic injury: perspectives in modern warfare. Ann Thorac Surg 2010;89(4):1035-1036

Reddick A, Ronald J, Morrison W: Intravenous fluid resuscitation: was Poiseuille right? Emerg Med J 2010;Epub ahead of print

Ritenour AE, Blackbourne LH, Kelly JF, et al: Incidence of primary blast injury in U.S. military Overseas Contingency Operations. Ann Surg 2010;251:1133-1137

Seamon M, Feather C, Smith B, et al: Just one drop: the significance of a single hypotensive blood pressure reading during trauma resuscitation. J Trauma 2010;68(6):1289-1295

Taylor DM, Bailey MS: A complication of the use of an intraosseous needle. JR Army Medical Corps 2010;156:132

Tintle S, Forsberg J, Keeling J, et al: Lower extremity combat-related amputations. J Surg Orthop Adv 2010;19(1):35-43

Van Exter: Prehospital endotracheal intubation in patients with severe traumatic brain injury: guidelines versus reality. Resuscitation 2010;81(5):627-628

Vandromme M, Griffin R, Weinburg J, et al: Lactate is a better predictor than systolic blood pressure for determining blood requirement and mortality: could prehospital measures improve trauma triage? J Am Coll Surg 2010;210(5):867-869

Vissers D, Stam W, Nolte T, et al: Efficacy of intranasal fentanyl spray versus other opioids for breakthrough pain in cancer. Curr Med Res Opin 2010;26(5):1037-1045

Vizcarra C, Clum S: Intraosseous route as alternative access for infusion therapy. J Infus Nurs 2010;33(3):162-174

Vlachou E, Gosling P, Moiemen N: Hydroxyethylstarch supplementation in burn resuscitation- a prospective randomized controlled trial. Burns 2010; Epub ahead of print

Walls R, Samuels-Kalow M, Perkins A: A new maneuver for endotracheal tube insertion during difficult glidescope intubation. J Emerg Med;39(1):86-88

Walsh K, Weant K, Cook A, et al: Potential benefits of high-dose methylprednisolone in acute spinal cord injuries. Orthopedics 2010;33(4):249-252

Wang H, Balasubramani G, Cook L, et al: Out-of-hospital endotracheal intubation experience and patient outcomes. Ann Emerg Med;55(6):527-537

Ward KR, Tiba H, Holbert WH, et al: Comparison of a new hemostatic agent to current combat hemostatic agents in a swine model of lethal extremity arterial hemorrhage. J Trauma 2007;63:276-284

## **May 2010**

Ahemd-Nusrath A, Nusrath M, Bryant D: Precautions required for needle cricothyroidotomy. *Br J Oral Maxillofac Surg* 2010;Epub ahead of print

Asensio J, Forno W, Castilla G, et al: Posterior ischemic optic neuropathy related to profound shock after penetrating trauma. *South Med J* 2002;95(9):1053-1057

Bassi M, Zuercher M, Erne J, et al: Endotracheal tube intracuff pressure during helicopter transport. *Ann Emerg Med* 2010;Epub ahead of print

Cheatham M, Safcsak K: Is the evolving management of intra-abdominal hypertension and abdominal compartment syndrome improving survival? *Crit Care Med* 2010;38(2):402-407

Cudnik M, Newgard C, Daya M, et al: The impact of rapid sequence intubation of trauma patient mortality in attempted prehospital intubation. *J Emerg Med* 2010;38(2):175-181

Demakis G, Rimland C: Untreated mild traumatic brain injury in a young adult population. *Arch Clin Neuropsychol* 2010;Epub ahead of print

Dente C, Shaz B, Nicholas J, et al: Early predictors of massive transfusion in patients sustaining torso gunshot wounded in a civilian level I trauma center. *J Trauma* 2010;68(2):298-304

Flaherty D, Hoxha B, Sun J, et al: Pyruvate-fortified fluid resuscitation improves hemodynamic stability while suppressing systemic inflammation and myocardial oxidative stress after hemorrhagic shock. *Mil Med* 2010;175(3):166

Fleishman R, Frazer D, Daya M, et al: Effectiveness and safety of fentanyl compared with morphine for out-of-hospital analgesia. *Prehosp Emerg Care* 2010;14(2):167-175

Hennessy D, Widder S, Zygun D, et al: Cervical spine clearance in obtunded blunt trauma patients: a prospective study. *J Trauma* 2010;68(3):576-582

Hillier R, Aboud A, Thind G, et al: Oral transmucosal fentanyl citrate: a novel analgesic agent for use in retinal photocoagulation. *Retina* 2009;29(10):1506-1512

Hink D, Franke A, Gatzka F: Use of vacuum-assisted closure negative pressure wound therapy in combat-related injuries-literature review. *Mil Med* 2010;175(3):173-181

Jeske H, Larndorfer R, Krappinger D, et al: Management of hemorrhage in severe pelvic injuries. *J Trauma* 2010;68(2):415-420

Kheirabadi B, Mace J, Terrazas I, et al: Safety evaluation of new hemostatic agents, smectite granules, and kaolin-combat gauze in a vascular injury wound model in swine. *J Trauma* 2010;68(2):269-278

Kandari J, Raizada S, Razzak A: Accidental laser injury to the eye. *Ophthalmic Surg Lasers Imaging* 2010;Epub ahead of print

Kudo D, Yamamura H, Nishiuchi T, et al: Anterior and posterior ischemic optic neuropathy related to massive fluid resuscitation after blunt trauma. *J Trauma* 2010;68(3):E67-E70

Lutes M, Worman D: An unanticipated complication of a novel approach to airway management. *J Emerg Med* 2010;38(2):222-224

Makley A, Goodman M, Friend L, et al: Resuscitation with fresh whole blood ameliorates the inflammatory response after hemorrhagic shock. *J Trauma* 2010;68(2):305-311

Mulligan R, Friedman J, Mahabir R: A nationwide review of the associations among cervical spine injuries, head injuries, and facial fractures. *J Trauma* 2010;68(3):587-592

Nelson B, MelnichE, Li J: Portable ultrasound for remote environments, part II: current indications. *J Emerg Med* 2010;Epub ahead of print

Newgard C, Schmicker R, Hedges J, et al: Emergency medical services intervals and survival in trauma: assessment of the “golden hour” in North American prospective cohort. *Ann of Emerg Med* 2010;55(3):235-246

Parkhouse D: Critical Care of military burn casualties at Role 3 facilities. *J R Army Med Corps* 2009;155(2):151-156

Petridis A, Doukas A, Barth H, et al: Outcome of craniocerebral gunshot injuries on the civilian population. Prognostic factors and treatment options. *Cen Eur Neurosurg* 2010: Epub ahead of print.

Salah N, Mhuircheartaigh R, Hayes N, et al: A comparison of four techniques of emergency transcricoid oxygenation in a manikin. *Anesth Analg* 2010;Epub ahead of print

Snell F, Halter M: A signature wound of war: mild traumatic brain injury. *J Psychosoc Nurs Ment Health Serv* 2010;48(2):22-28

Takasu A, Minagawa Y, Ando S, et al: Improved survival time with combined early blood transfusion and fluid administration in uncontrolled hemorrhagic shock in rats. *J Trauma* 2010;68(2):312-316

Tong W, Zheng P, Xu J, et al: Early CT signs of progressive hemorrhagic injury following acute traumatic brain injury. *Neuroradiology* 2010;

Wang H, Balasubramani G, Cook L, et al: Out-of hospital endotracheal intubation experience and patient outcomes. *Ann Emerg Med* 2010;Epub ahead of print.

Weintrob A, Roediger M, Barber M, et al: Natural history of colonization with gram-negative multidrug-resistant organisms among hospitalized patients. *Infect Control Hosp Epidemiol* 2010;31(4):330-337

Zugai B, Eley V, Malitt K, et al: Practice patterns for predicted difficult airway management and access to airway equipment by anaesthetists in Queensland, Australia. *Anaesth Intensive Care* 2010;38(1):27-32

## February 2010

Abernathy J, Reeves S: Airway catastrophes. Curr Opin Anaesthesiol 2010;23(1):41-46

Belmont P, Goodman G, Zacchilli M, et al: Incidence and epidemiology of combat injuries sustained during “the surge” portion of operation Iraqi Freedom by a U.S. Army brigade combat team. J Trauma 2010;68(1):204-210

Bouillon B, Brohi K, Hess J, et al: Educational initiative on critical bleeding in trauma: Chicago, July 11-13, 2008. J Trauma 2010;68(1):225-230

Burgert J: Intraosseous infusion of blood products and epinephrine in an adult patient in hemorrhagic shock. AANA J 2009;77(5):359-363

Dong W, Cai B, Peña G, et al: Ethyl pyruvate prevents inflammatory responses and organ damage during in porcine hemorrhage. Shock 2009;Epub ahead of print

Eastridge B, Costanzo G, Jenkins D, et al: Impact of joint theater trauma system initiatives on battlefield injury outcomes. Am J Surg 2010;198(6):852-857

Flint N, Russell W, Thompson J: Comparison of different methods of ventilation via cannula cricothyroidotomy in a trachea-lung model. Br J Anaesth 2009;103(6):891-895

Franschman G, Greuters S, Loer S, et al: Prehospital treatment guidelines in severe traumatic brain injury: What really happens outside the hospital? Resuscitation 2009;Epub ahead of print.

Goon S, Stephens R, Smith H: The emergency airway. Br J Hosp Med 2010;70(12):M186-188

Hargreaves S, Lacasia C, Groom P: Regional audit- availability of equipment and training in emergency cricothyroidotomy. Anaesthesia 2010;65(1):107

Haut E, Kalish B, Efron D, et al: Spine immobilization in penetrating trauma: more harm than good? J Trauma 2010;68(1):115-121

Hetzler M, Rick G: Damage control resuscitation for the special forces medic-simplifying and improving prolonged trauma care-part two. J Spec Oper Med 2009;9(4):53-62

Holcomb J, Spinella P: Optimal use of blood in trauma patients. *Biologicals* 2010; Epub ahead of print

Hoyt B: Blood and war-lest we forget. *J Am Coll Surg* 2009;209(6):681-6

Jiao W, Kiang J, Cary L, et al: COX-2 inhibitors are contraindicated for treatment of combined injury. *Radiat Res* 2009;172(6):686-697

Lamhaut L, Dagron C, Apriotesei R, et al: Comparison of intravenous and intraosseous access by pre-hospital medical emergency personnel with and without CBRN protective equipment. *Resuscitation* 2009; Epub ahead of print

Lee J, Kim K, Jo Y, et al: Early norepinephrine infusion delays cardiac arrest after hemorrhagic shock in rats. *J Emerg Med* 2009;37(4):376-382

Lew T, Walker J, Wenke J, et al: Characterization of craniomaxillofacial battle injuries sustained by the United States service members in the current conflicts of Iraq and Afghanistan. *J Oral Maxillofac Surg* 2010;68(1):3-7

Ling G, Rhee P, Ecklund J: Surgical innovations arising from the Iraq and Afghanistan wars. *Annu Rev Med* 2010;61:457-468

Morrison J, Mountain A, Galbraith K, et al: Penetrating pelvic battlefield trauma; internal use of chitosan-based haemostatic dressings. *Injury* 2010;41(2):239-241

Napolitano L, Kurek S, Luchette F, et al: Clinical practice guideline: red blood cell transfusion in adult trauma and critical care. *Crit Care Med* 2009;37(12):3124-3157

Nelson B, Melnick E, Li J: Portable ultrasound for remote environments, part i: feasibility of field deployment. *J Emerg Med* 2010;Epub ahead of print

Rockswold G, Solid C, Paredes-Andrade E, et al: Hypertonic saline and its effect on intracranial pressure, cerebral perfusion pressure, and brain tissue oxygen. *Neurosurgery* 2009;65(6):1035-1041

Sánchez-Morillas L, Rojas Pérez-Ezquerra P, Reaño-Martos M, et al: Systemic anaphylaxis caused by moxifloxacin. *Allergol Immunopathol* 2010;Epub ahead of print

Schuster K, Lofthouse R, Moore C, et al: Pulseless electrical activity, focused abdominal sonography for trauma, and cardiac contractile activity as predictors of survival after trauma. *J Trauma* 2009;67(6):1154-1157

Sen A, Gulati A: Use of Magnesium in traumatic brain injury. Neurotherapeutics 2010;7(1):91-99

Sharp C, Mangram A, Lorenzo M, et al: A major metropolitan “field amputation” team: a call to arms ... and legs. J Trauma 2009;67(6):1158-1161

Sondén A, Rocksén D, Riddez L, et al: Trauma attenuating backing improves protection against behind armor blunt trauma. J Trauma 2009;67(6):1191-1199

Ward K, Archambault R, Mersfelder T: Severe adverse skin reaction to nonsteroidal anti-inflammatory drugs: a review of literature. Am J Health Syst Pharm 2010;67(3):206-213

## **November 2009**

Alam H, Bice L, Butt M, et al: Testing of blood products in a polytrauma model: results of a multi-institutional randomized preclinical trial. *J Trauma* 2009;67(4):856-864

Allard C, Scarpelini S, Rhind S, Baker A, et al: Abnormal coagulation test are associated with progression of traumatic intracranial hemorrhage. *J Trauma* 2009;67(5):959-967

Ambrus R, Kocbek P, Kristi J, et al: Investigation of preoparation parameters to improve the dissolution of poorly water-soluble meloxicam. *Int J Pharm* 2009;381(2):153-159

Arnaud F, Teranishi K, Tomori T, et al: Comparison of 10 hemostatic dressing in a groin puncture model in swine. *J Vasc Surg* 2009;50(3):632-639

Arnaud F, Parrento-Sadalan D, Tomori T, et al: Comparison of 10 hemostatic dressings in a groin transaction model in swine. *J Trauma* 2009;67:848-855

Batista M, Bravin A, Lopes L, et al: Pressor response to fluid resuscitation in endotoxic shock: involvement of vasopressin. *Crit Care Med* 2009;37(11):2968-2972

Blackwell T, Kline J, Willis J, et al: Lack of association between prehospital response times and patient outcomes. *Prehosp Emerg Care* 2009;13:444-450

Buckenmaier C, Rupprecht C, McKnight G, et al: Pain following battlefield injury and evacuation: a survey of 110 casualties for the wars in Iraq and Afghanistan. *Pain Med* 2009;Oct: Epub ahead of print

Cabralles P: Low dose nitrite enhances perfusion after fluid resuscitation from hemorrhagic shock. *Resuscitation* 2009;Oct:Epub ahead of print

Cheatham M, Safcsak K: Is the evolving management of intra-abdominal hypertension and abdominal compartment syndrome improving survival? *Crit Care Med* 2009;Sept;Epub ahead of print

Cho R, Bakken H, Reynolds M, et al: Concomitant cranial and ocular combat injuries during Operations Iraq Freedom. *J Trauma* 2009;67(3):5126-520

Chung K, Wolf S, Cancio L, et al: Resuscitation of severely burned military casualties; fluid begets more fluid. *J Trauma* 2009;67(2):231-237

Cobas M, De la Peña M, Manning R et al: Prehospital intubations and mortality: a level 1 trauma center perspective. *Anesth Analg* 2009; 109(2):489-493

Cotton B, Jerome R, Collier B, et al: Guidelines for prehospital fluid resuscitation in the injured patient. *J Trauma* 2009;67(2):389-402

Elliot M, Jallo J, Barbe M, et al: Hypertonic saline attenuates tissue loss and astrocyte hypertrophy in a model of traumatic brain injury. *Brain Res* 2009:Epub ahead of print

Fatovich D, Jacobs I: The relationship between remoteness and trauma deaths in Western Australia. *J Trauma* 2009;67(5):910-914

Flint N, Russell W, Thompson J: Comparison of different methods of ventilation via cannula cricothyroidotomy in a trachea-lung model. *Br J Anaesth* 2009;Sept:Epub ahead of print.

Fludger S, Bell A: Tourniquet application in a rural Queensland HEMS environment. *Air Med J* 2009;28(6):291-293

Franschman G, Peerdeman S, Greuters S, et al: Prehospital endotracheal intubation in patients with severe traumatic brain injury: guidelines verses reality. *Resuscitation* 2009;80(10):1147-1151

Gerhardt R, Matthews J, Sullivan S: The effect of systemic antibiotic prophylaxis and wound irrigation on penetrating combat wounds in a return-to-duty population. *Prehosp Emerg Care* 2009;13(4):500-504

Gonzalez C, Brunstein A: Training for emergencies. *J Trauma* 2009;67(2):S100-105

Greenburg A: The ideal blood substitute. *Crit Care* 2009;25:415-424

Hardaway R: One surgeon's army experience with "wound shock" from Pearl Harbor to present. *Mil Med* 2009;174(9):944-947

Hecht-Dolnick M, Barkan H, Taharka A, et al: Hetastarch increases the risk of bleeding complications in patients after off-pump coronary bypass surgery: a randomized clinical trial. *J Thorac Cardiovasc Surg* 2009;138(3):703-711

Holmes J, Wisner D, McGahan J, et al: Clinical prediction rules for identifying adults at very low risk for intra-abdominal injuries after blunt trauma. *Annals of Emerg Med*;54(4):575-584

Hospital del Henares, Madrid, Spain: Steroid use in acute lung injury. *Crit Care Med* 2009;37(9):2680

Johansson P, Stissing T, Bochsen L, et al: Thrombelastography in assessing coagulopathy in trauma. Scan J Trauma Resusc Emerg Med 2009;17(1):45

Kaspar R, Griffith M, Mann P, et al: Association of bacterial colonization at the time of presentation to a combat support hospital in a combat zone with subsequent 30-day colonization or infection. Mil Med 2009;174:899-903

Kattail D, Furlan J, Fehlings M: Epidemiology and clinical outcomes of acute spine trauma and spinal cord injury: experience from a specialized spine trauma center in Canada in comparison with a large national registry . J Trauma 2009;67(5):936-943

Kheirabadi B, Scherer M, Estep J, et al: Determination of efficacy of new hemostatic dressings in a model of extremity arterial hemorrhage in swine. J Trauma 2009;67(3):459-460

Kiraly L, Underwood S, Differding J, et al: Transfusion of aged packed red blood cells results in decreased tissue oxygenation n critically injured trauma patient. J Trauma 2009;67(1):29-32

Knapp B, Tsuchitani S, Sheele J, et al: Prospective evaluation of an emergency medical services-administered alternative transport protocol. Prehosp Emerg Care 2009;13:432-436

Leidel B, Kirchhoff C, Bogner V, et al: Is there intraosseous access route fast and efficacious compared to conventional central venous catheterization in adult patients under resuscitation in the emergency department? A prospective study observational pilot study. Patient Saf Surgery 2009;8:24

Levitan R, Bortle C, Snyder T, et al: Use of a battery operated needle driver for intraosseous access by novice users: skill acquisition with cadavers. Annals of Emerg Med 2009;54(5): 692-694

Mahambrey T, Fowler A, Pinto R, et al: Early massive transfusion in trauma patients: Canadian single-centre retrospective cohort study. Can J Anaesth 2009;56(10):740-750

McQuay N, Cipolla J, Franges E, et al: The use of recombinant activated factor VIIa in coagulopathic traumatic brain injury requiring emergent craniotomy: is it beneficial? J Neurosurg 2009;111(4):666-671

Menon D, Zahed C: Prediction of outcome in severe traumatic brain injury. Curr Opin Crit Care 2009;15(5):437-441

Metzger J, Eastman A, Benitez, et al: The livesaving potential of specialized on-scene medical support for urban tactical operations. *Prehosp Emerg Care* 2009;13(4):528-531

Mistry N, Bleetman A, Roberts K: Chest decompression during the resuscitation of patients in prehospital traumatic cardiac arrest. *Emerg Med J* 2009;26(10):738-740

Nishijima D, Zehtabchi S: The efficacy of recombinant activated Factor VIIa in severe trauma. *Annals of Emerg Med* 2009;54(5):737-744

Parker P: Emergency Tourniquet use. *Ann Surg* 2009;250(3):496-497

Paxton J, Knuth T, Klausnert H: Proximal humerus intraosseous infusion: a preferred emergency venous access. *J Trauma* 2009;67(3):606-611

Ragal B, Allred C, Brevard S, et al: Fracture of the thoracolumbar spine sustained by soliders in vehicles attacked by improvised explosive devices. *Spine* 2009: Epub ahead of print.

Sarkar D, Demma A, Stulz D, et al: Expect the unexpected: two cases of penetrating head and neck trauma for Operation Iraqi Freedom. *Ear Nose Throat J* 2009;88(9):E19-21

Shavit I, Hoffmann Y, Galbraith R, et al: Comparison of two mechanical intraosseous infusion devices: a pilot, randomized crossover trial. *Resuscitation* 2009;80(9):1029-1033

Solomkin J, Zhao Y, Ma E, et al: Moxifloxacin is non-inferior to combination therapy with ceftriaxone plus metronidazole in patients with community-origin complicated intra-abdominal infections. *Int J Antimicrob Agents* 2009: Epub ahead of print

Stansbury L, Dutton R, Stein D, et al: Controversy in trauma resuscitation: do ratios of plasma to red blood cells matter? *Transfusion Med Rev* 2009;23(4):255-265

Sztajnkrycer M: Needle thoracostomy by non-medical law enforcement personnel: preliminary data on knowledge retention. *Prehosp Disaster Med* 2009;23(6):553-557

Thomas R, McManus J, Johnson A, et al: Ocular injury reduction for ocular protection use in current combat operations. *J Trauma* 2009;66:S99-1032

Towner K: Acinetobacter: an old friend, but a new enemy. *J Hosp Infect* 2009;Epub ahead of print

Vick-Fragoso R, Hernández-Oliva G, Cruz-Alcázar J, et al: Efficacy and safety sequential intravenous/oral Moxifloxacin vs intravenous/oral Amoxicillin/Clavulanate for complicated skin and skin structure infections. Infections;Epub ahead of print.

Von Elm E, Schoettker P, Henzi I, et al: Pre-hospital tracheal intubation in patients with traumatic brain injury; systematic review of current evidence. Br J Anaesth 2009;103(3):371-386

Wafaisade A, Lefering R, Tjardes T, et al: Acute coagulopathy in isolated blunt traumatic brain injury. Neurocrit Care 2009;Epub ahead of print.

Yaghoubian A, Lewis R, Putnam B, et al: Reanalysis of prehospital intravenous fluid administration in patients with penetrating truncal injury and field hypertension. Am Surg 2009;73(10):1027-1030

Zhang R, Hunter R, Gonzalez E, et al: Poloxamer 188 prolongs survival of hypotensive resuscitation and decreases vital tissue injury after full resuscitation. Shock 2009;32(4):442-450

## **August 2009**

Allison C, Trunkey D: Battlefield trauma, traumatic shock and consequences: war-related advances in critical care. Crit Care Clin 2009;25:31-45

Beilman G, Blondet J, Nelson T, et al: Early hypothermia in severely injured trauma patients is a significant risk factor for multiple organ dysfunction syndrome but not mortality. Ann Surgery 2009;249:845-850

Brenner T, Gries A, Helm M, et al: Intraosseous infusion systems in the prehospital setting. Resuscitation 2009;80:607

Brown M, Daya M, Worley J, et al: Experience with chitosan dressing in a civilian EMS system. J Emerg Med 2009;37:1-7

Cai B, Deitch E, Grande D, et al: Anti-inflammatory resuscitation improves survival in hemorrhage with trauma. J Trauma 2009;66:1632-1640

Carter A, Davis K, Evans L, et al: Information loss in emergency medical services handover trauma patients. Prehosp Emerg Care 2009;13:280-285

Champion H, Holcomb J, Young L: Injuries from explosions: physics, biophysics, pathology, and required research focus. J Trauma 2009;66:1468-1477

Chen L, Resiner A, Gribok A, et al: Exploration of prehospital vital sign trends for the prediction of trauma outcomes. Prehosp Emerg Care 2009;13:286-294

Cobas M, De la Peña M, Manning R, et al: Prehospital intubations and mortality: a level 1 trauma center perspective. Anesth Analg 2009;109:489-493

Cooper E, Bateman S, Muir W: Evaluation of hyperviscous fluid resuscitation in a canine model of hemorrhagic shock: a randomized, controlled study.

Denver Metro Airway Study Group: A prospective multicenter evaluation of prehospital airway management performance in a large metropolitan region. Prehosp Emerg Care 2009;13:304-310

Dougherty A, Mohrie C, Galarneau M, et al: Battlefield extremity injuries in Operation Iraq Freedom. Injury 2009;40:772-777

Ford N, Mills E, Zachariah R, et al: Ethics of conducting research I conflict settings. Confl Health 2009;10:3-7

Gauna AA, Kang SK, Triano ML, et al: Oral Transmucosal fentanyl citrate for dyspnea in terminally ill patients: an observational case series. *J Palliat Med* 2009;11:643-648

Gerritse B, Scheffer G, Draisma J: Prehospital intraosseous access with the bone injection gun by a helicopter-transported emergency medical team. *J Trauma* 2009;66:1739-1741

Huang C, Kao K, Hsu K, et al: Effects of hydroxyethyl starch resuscitation on extravascular lung water and pulmonary permeability in sepsis-related acute respiratory distress syndrome. *Crit Care Med* 2009;37:1948-1955

Johannigman J, Branson R, Edwards M: Closed loop control of inspired oxygen concentration in trauma patients. *J Amer Col of Surgeon* 2009;208:763-768

Kilbourne M: Hemostatic efficacy of modified amylopectin powder in a lethal porcine model of extremity arterial injury. *Ann Emerg Med* 2009;53:804-810

Kucewicz J, Bailey M, Kaczkowski P, et al: Focused ultrasound: concept for automated transcutaneous control of hemorrhage in austere settings. *Aviat Space Environ Med*;2009;80:391-394

Lam N, Hagberg C, Bassili L: Use of the video laryngoscopy for combitube exchange in a difficult airway. *J Clin Anesth* 2009;21:294-296

Langeron O, Birenbaum A, Amour J: Airway management in trauma. *Minerva Anestesiol* 2009;75:307-311

Levitan R, Bortle C, Snyder T, et al: Use of a battery-operated needle driver for intraosseous access by novice users: skill acquisition with cadavers. *Ann Emerg Med* 2009;Epub ahead of print.

Luck R, Haines C, Mull C: Intraosseous access. *J Emerg Med* 2009;Epub ahead of print

Lundgren J, Henriksson O, Pretorius T, et al: Field torso-warming modalities: a comparative study using a human model. *Prehops Emerg Care* 2009;13:371-378

Manring M, Hawk A, Calhoun J, et al: Treatment of war wounds: a historical review. *Clin Orthop Relat Res* 2009;Epub ahead of print

Martínez-Odriozola P, Gutiérrez-Macías A, Ibarmia-Lahuerta J, et al: Meloxicam as a cause of drug-induced autoimmune hepatitis. *Dig Dis Sci* 2009;Epub of print

McArdle G, McAuley D, McKinley A, et al: Preliminary results of a prospective randomized trial of restrictive versus standard fluid regime in elective open abdominal aortic aneurysm repair. Ann of Surgery 2009;Epub ahead of print.

McQuay N, Cipolla J, Franges E, et al: The use of recombinant activated factor VIIa in coagulopathic traumatic brain injuries requiring emergent craniotomy: is it beneficial? J Neurosurg 2009;Epub ahead of print

Mullins J, Harrahill M: Use of a tourniquet after a gunshot wound to the thigh. J Emerg Nurs 2009;35:265-267

Neparidze N, Kriz N, Ornstein D: Hetastarch haemophilia . Haemophilia 2009;15:976-979

Otero V, Posse J, Carmona I, et al: Efficacy of fluoroquinolones against pathogenic oral bacteria. Mini Rev Med Chem 2009:Epub ahead of print.

Plurad D, Chandrasoma S, Best C, et al: A complication of intracorporeal use of quickclot for pelvic hemorrhage. J Trauma 2009;66:1482-1484

Ramasamy A, Harrisson S, Lasrado I, et al: A review of casualties during the Iraq insurgency 2006- a British field hospital experience. Injury 2009;40:493-497

Ramasamy A, Midwinter M, Mahoney P, et al: Learning the lessons from conflict: Pre-hospital cervical spine stabilization following ballistic neck trauma. Injury 2009:Epub ahead of print

Sambasivan C, Cho S, Zink K, et al: A highly porous silica and chitosan-based hemostatic dressing is superior in controlling hemorrhage in a severe groin injury model in swine. Am J Surgery 2009;197:576-580

Schmid M, Mang H, Ey K, et al: Prehospital airway management on rescue helicopters in the United Kingdom. Anaesthesia 2009;64:625-631

Streets C: Lessons from the battlefield in the management of major trauma. Br J Surg 2009;96:831-832

Tam R, Maloney J, Gaboury I, et al: Review of endotracheal intubations by Ottawa Advanced Care Paramedics in Canada. Prehosp Emerg Care 2009;13:311-315

Thaveau F, Zoll J, Bouitbir J, et al: Contralateral leg as a control during skeletal muscle ischemia-reperfusion. J Surg Res 2009;155:65-69

Wang H, Weaver M, Abo B, et al: Ambulance stretcher adverse events. Qual Saf Health Care 2009;18:213-216



## **April 2009**

Archam S, Prause G, Gumpert R, Seibert FJ, Kügler B: Cricothyroidotomy on the scene in a patient with severe facial trauma and difficult neck anatomy. *AM J Emerg Med* 2009;27:133

Bauman RA, Ling GS, Tong L, et al: An introductory characterization of a combat casualty care-relevant swine model of closed head injury resulting from exposure to explosive blast. *J Neurotrauma* 2009 Feb 12; Epub ahead of print

Black IH, McManus J: Pain Management in current combat operations. *Prehosp Emerg Care* 2009;13:223-227

Bledsoe B, Gandy W: The Disappearing Endotracheal Tube: Historic skill threatened by lack of practice and new devices. *JEMS* 2009;34:88-99

Bruns B, Gentillello L, Elliot A, Shafi S: Prehospital hypotension redefined. *J Trauma* 2008;65:1217-1221

Bullard-Berent J: Intraosseous infusion: an instructional program for healthcare providers, revised edition (CD-ROM) *Prehosp Emerg Care* 2009;13:266

Cai B, Chen F, Lin X, et al: Anti-inflammatory adjuvant in resuscitation fluids improves survival in hemorrhage: *Crit Care Med* 2009;37:1133-1134

Chan SS: Tension pneumothorax managed without immediate needle decompression. *J Emerg Med* 2009;36:242-245

David JS, Dubien PY, Capel O, Peguet O, Gueugniaud PY: Intraosseous infusion using the bone injection gun in the prehospital setting. *Resuscitation* 2009;80:384-385

Devlin JJ, Kircher S, Kozen BG, Littlejohn LF, Johnson AS: Comparison of ChitoFlex®, CELOX, and QuikClot® in Control Hemorrhage. *J Emerg Med* 2009; Apr 1 - Epub ahead of print

Fatovich DM, Jacobs IG: The relationship between remoteness and trauma deaths in Western Australia. *J Trauma* 2008 Aug 15 - Epub ahead of print

Gehardt RT, De Lorenzo RA, Oliver J, Holcomb JB, Pfaff JA: Out-of-hospital combat casualty care in the current war in Iraq. *Ann Emerg Med* 2009;53:169-172

Hartog C, Brunkhorst F, Reinhart K: Hydroxyethyl starch 130/0.4 and surgical blood loss. *Anesth Analg* 2009;108:672

Hasan SM, Shoaib MH, Hassan F, Rehman I: Bioequivalence studies of two brands of meloxicam tablets in healthy Pakistani volunteers. *Pak J Pharm Sci* 2009;22:199-204

Hassis VJ, Schino MA, Tepas JJ, et al: Definitive establishment of airway control is critical for optimal outcome in lower cervical spinal cord injury. *J Trauma* 2008;65:1328-1332

Holcomb JB, McManus JG, Kerr ST, Pusateri AE: Needle versus tube thoracostomy in a swine model of traumatic tension hemopneumothorax. *Prehosp Emerg Care.* 2009;13:18-27

Hurtado TR, Montoya C: Trauma scissors vs. the rescue hook, exposing a simulated patient: a pilot study. *J Emerg Med* 2009;36:232-235

Johannigman JA, Branson R, Lecroy D, Beck G: Autonomous control of inspired oxygen concentration during mechanical ventilation of the critically injured trauma patient. *J Trauma* 2009;66:386-392

Kheirabadi BS, Edens JW, Terrazas IB et al: Comparison of new hemostatic granules/powders with currently deployed hemostatic products in a lethal model of extremity arterial hemorrhage in swine. *J Trauma* 2009;66:316-328

Kragh JF Jr, Walters TJ, Baer DG, Fox CJ, Wade CE, Salinas J, Holcomb JB: Survival with emergency tourniquet use to stop bleeding in major limb trauma. *Ann Surg* 2009;249:1-7

Lacquiere A, Heard AM: Emergency cricothyroidotomy: training is paramount and oxygenation is the aim. *Anaesthesia* 2009;64:447-448

Moore FA: Tourniquets: Another adjunct in damage control? *Ann Surg* 2009;249:8-9

Parker PJ: Pre-Hospital Antibiotic Administration. *JR Army Med Corps* 2009;154:5-9

Schramko AA, Suojaranta-Ylinen RT, Kuitunen AH, Kukkonen SI, Niemi TT: Rapidly degradable hydroxyethyl starch solutions impair blood coagulation after cardiac surgery: a prospective randomized trial. *Anesth Analg* 2009;108:30-36

Shipman N, Lessard CS: Pressure applied by the emergency Israeli bandage. *Mil Med* 2009;174:86-92

Sise MJ, Shackford SR, Sise CB, et al: Early intubation in the management of trauma patients; indications and outcomes in 1,000 consecutive patients. *J Trauma* 2009;66:32-9

Spinella PC, Perkins JG, Grathwohl KW, Beekley AC, Holcomb JB: Warm fresh whole blood is independently associated with improved survival for patients with combat-related traumatic injuries. *J Trauma* 2009;66:S69-S76

Stein DM, Dutton RP, Kramer ME, Scalea TM: Reversal of coagulopathy in critically ill patients with traumatic brain injury: recombinant factor VIIa is more cost-effective than plasma. *J Trauma* 2009;66:63-72

Stevens RL, Rochester AA, Busko J, et al: Needle thoracostomy for tension pneumothorax: failure predicted by chest computed tomography. *Prehosp Emerg Care* 2009;13:14-17

Talving P, Benfield R, Hadjizachari P, et al: Coagulopathy in severe traumatic brain injury: a prospective study. *J Trauma* 2009;66:55-61

Thistlethwaite R, Ray R, Kessell G: Emergency cricothyroidotomy: equipment in anaesthetic practice. *Anaesthesia* 2009;64:448-449

Velmahos GC, Tabbara M, Spaniolas K, et al: Self-expanding hemostatic polymer for control of exsanguinating extremity bleeding. *J Trauma* 2009;66:984-8

Warner KJ, Sharar SR, Copass MK, Bulger EM: Prehospital management of the difficult airway: a prospective cohort study. *J Emerg Med* 2009;36:257-265

## **February 2009**

Cernal L: Emergency medicine physicians highlight “Army-wide emergency medical care deficiency” in training of flight medics. Ann Emerg Med 2008;52:522-524

Gerhardt RT, DeLorenzo RA, Olivier J, Holcomb JB, Pfaff JA: Out-of-hospital combat casualty care in the current war in Iraq. Ann Emerg Med 2009;53:169-174

Kragh JF, Walters TJ, Baer, DJ, et al: Survival with emergency tourniquet use to stop bleeding in major limb trauma. Ann Surg 2009;249:1-7

Shipman N, Lessard CS: Pressure applied by the Emergency/Israeli bandage. Milit Med 2009;174:86-92

Suter RE: Forward medical care and the military emergency medicine workforce: too much demand and not enough supply? Ann Emerg Med 2009;53:175-177

Wamer KJ, Cuschieri J, Garland B, et al: The utility of early end-tidal capnography in monitoring ventilation status after severe injury. J Trauma 2009;66:26-31

## **November 2008**

Brodie S, Hodgetts TJ, Ollerton J, et al: Tourniquet use in combat trauma: UK military experience. *J R Army Med Corps* 2007;153:310-313

Calhoun JH, Murray CK, Manring MM: Multidrug-resistant organisms in military wounds from Iraq and Afghanistan. *Clin Orthop Relat Res* 2008;466:1356-1362

Englehart MS, Cho SD, Tieu BH, et al: A novel highly porous silica and chitosan-based hemostatic dressing is superior to HemCon and gauze sponges. *J Trauma* 2008;65:890-892

Hess JR, Brohi K, Dutton RP, et al: The coagulopathy of trauma: a review of mechanisms. *J Trauma* 2008;65:748-754

Holcomb JB, Wade CE, Michalek JE, et al: Improved plasma and platelet to red blood cell ratios improves outcome in 466 massively transfused civilian trauma patients. *Ann Surg* 2008;248:in press

Ivascu FA, Howells GA, Junn FS, et al: Predictors of mortality in trauma patients with intracranial hemorrhage on preinjury aspirin or clopidogrel. *J Trauma* 2008;65:785-788

Kalish J, Burke P, Feldman J, et al: The return of tourniquets. *JEMS* 2008;33:44-46

Kozek-Langenecker SA, Jungheinrich C, Sauermann W, Van der Linden P: The effects of hydroxyethyl starch 130/0.4 (6%) on blood loss and use of blood products in major surgery: a pooled analysis of randomized clinical trials. *Anesth Analg* 2008;107:382-390

Lier H, Krep H, Schroeder S, Stuber F: Preconditions of hemostasis in trauma: a review. The influence of acidosis, hypocalcemia, anemia, and hypothermia on functional hemostasis in trauma. *J Trauma* 2008;65:951-960

Moskowitz DM, Shander A, Javidroozi M, et al: Postoperative blood loss and transfusion associated with use of Hextend in cardiac surgery patients at a blood conservation center. *Transfusion* 2008;48:768-775

Nichol G, Thomas E, Callaway, et al: Regional variation in out-of-hospital cardiac arrest incidence and outcome. *JAMA* 2008;300:1423-1431

Rathinam S, Beetman A, Steyn RS: Needle thoracostomy in treatment of a tension pneumothorax. *J Trauma* 2008;65:964

Sasson C, Hegg AJ, Macy M, et al: Prehospital termination of resuscitation in cases of refractory out-of-hospital cardiac arrest. JAMA 2008;300:1432-1438

Stiver SI, Manley GT: Prehospital management of traumatic brain injury. Neurosurg Focus 2008;25:E5

Tien HC, Jung V, Rizoli SB, Acharya SV, MacDonald JC: An evaluation of Tactical Combat Casualty Care interventions in a combat environment. J Am Coll Surg 2008;207:174-178

Weeks DL, Jahr JS, Lim JC, Butch AW, Driessen B: Does Hextend impair coagulation compared to 6% hetastarch? An ex-vivo thromboelastography study. Am J Ther 2008;15:225-230

## August 2008

Adams BD, Cuniowski PA, Muck A, De Lorenzo RA: Registry of emergency airways arriving at combat hospitals. *J Trauma* 2008;64:1548-1554

Beekley AC: Damage control resuscitation: a sensible approach to the exsanguinating surgical patient. *Crit Care Med* 2008;36:S267-S274

Beninati W, Meyer MT, Carter TE: The critical care air transport program. *Crit Care Med* 2008;36:S370-S376

Blackbourne LH: Combat damage control surgery. *Crit Care Med* 2008;36:S304-S310

Brethauer SA, Chao A, Chambers LW, Green DJ, Brown C, Rhee P, Bohman HR: Invasion vs Insurgency: US Navy/Marine Corps Forward Surgical Care During Operation Iraqi Freedom. *Arch Surg.* 2008;143(6):564-569.

Dewitt DS, Prough DS: Blast-induced brain injury and post-traumatic hypotension and hypoxemia. *J Neurotrauma* 2008: E-pub ahead of print 30 April

Doyle GS, Taillac PP: Tourniquets: A review of current use with proposals for expanded prehospital use. *Prehospital Emergency Care* 2008;12:241-256

Fergusson DA, McIntyre L: The future of clinical trials evaluating blood substitutes. *JAMA* e-pub 28 April 2008

Kheirabadi BS, Crissey JM, Deguzman R, et al: Effects of synthetic versus natural colloid resuscitation on inducing dilutional coagulopathy and increasing hemorrhage in rabbit. *J Trauma* 2008;64:1218-1228

Johannigman JA: Maintaining the continuum of en route care. *Crit Care Med* 2008;36:S377-S382

Kortbeek JB, et al: Advanced Trauma Life Support, 8<sup>th</sup> Edition, The evidence for change. *J Trauma* 2008;64:1638-1650

Kozek-Langenecker SA, Jungheinrich C, Sauermann W, Van der Linden P: The effects of hydroxyethyl starch 130/0.4 (6%) on blood loss and use of blood products in major surgery: a pooled analysis of randomized clinical trials. *Anesth Anal* 2008;107:382-390

Mabry R, McManus JG: Prehospital advances in the management of severe penetrating trauma. *Crit Care Med* 2008;36:S258-266

Malchow RJ, Black IH: The evolution of pain management in the critically ill trauma patient: emerging concepts from the global war on terrorism. Crit Care Med 2008;36:S346-S347

McIntosh SE, Swanson ER, Barton ED: Cricothyroidotomy in air medical transport. J Trauma 2008;64:1543-1547

Murray CK: Infectious disease complications of combat-related injuries. Crit Care Med 2008;36:S358-S364

Natanson C, Kern SJ, Lurie P, Banks SM, Wolfe SM: Cell-free hemoglobin-based substitutes and risk of myocardial infarction and death. JAMA e-pub 28 April 2008

Niles SE, McLaughlin DF, Perkins JG, Wade CE, Li Y, Spinella PC, Holcomb JB: Increased mortality associated with the early coagulopathy of trauma in combat casualties. J Trauma 2008;64:1459-1463

Perkins JG, Cap AP, Weiss BM, Reid TJ, Bolan CE: Massive transfusion and nonsurgical hemostatic agents. Crit Care Med 2008;36:S325-S329

Richey SL: Tourniquets for the control of traumatic hemorrhage: a review of the literature. World Journal of Emergency Surgery 2006

## April 2008

Beekley AC, Sebesta JA, Blackbourne LH, Herbert GS, Kauvar DS, Baer DG, Walters TJ, Mullenix PS, Holcomb JB: Prehospital tourniquet use in Operation Iraqi Freedom: effect on hemorrhage control and outcomes. *J Trauma* 2008;64:S28-S37

Calhoun JH, Murray CK, Manring MM: Multidrug-resistant organisms in military wounds from Iraq and Afghanistan. *Clin Orthop Relat Res* 2008; E-pub ahead of print

Cancio LC, Wade CE, West SA, Holcomb JB: Prediction of mortality and of the need for massive transfusion in casualties arriving at combat support hospitals in Iraq. *J Trauma* 2008;64:S51-S55

Conger NG, Landrum ML, Jenkins DH, et al: Prevention and management of infections associated with combat-related thoracic and abdominal cavity injuries. *J Trauma* 2008;64:S257-S264

Convertino VA, Ryan KL, Rickards CA, Salinas J, McManus JG, Cooke WH, Holcomb JB: Physiological and medical monitoring for en route care of combat casualties. *J Trauma* 2008;64:S342-S353

Cordts PR, Brosch LA, Holcomb JB: Now and then: combat casualty care policies for Operation Iraqi Freedom and Operation Enduring Freedom compared with those of Vietnam. *J Trauma* 2008;64:S14-S20

Dayan L, Zinmann C, Stahl S, Norman D: Complications associated with prolonged tourniquet application on the battlefield. *Milit Med* 2008;173:63-66

Doyle GS, Taillac PP: Tourniquets – a review of current use with proposals for expanded prehospital use. *Prehosp Emerg Care* 2008;12:241-256

Hospenthal DR, Murray CK, Anderson RC, et al: Guidelines for the prevention of infection after combat-related injuries. *J Trauma* 2008;64:S211-S220

Kelly JF, Ritenhour AE, McLaughlin DF, Bagg KA, Apodaca AN, Mallak CT, Pearse L, Lawnick MM, Champion HR, Holcomb JB: Injury severity and causes of death from Operation Iraqi Freedom and Operation Enduring Freedom: 2003-2004 vs 2006. *J Trauma* 2008;64:S21-S26

Kragh JF, Walters TJ, Baer DG, Fox CJ, Wade CE, Salinas J, Holcomb JB: Practical use of emergency tourniquets to stop bleeding in major limb trauma. *J Trauma* 2008;64:S38-S50

Lott MN, Fuller JJ, Hancock HA, Singh J, McGwin G, Marcus DM: Vitreal penetration of oral moxifloxacin in humans. *Retina* 2008;28:473-476

Madigan MC, Kemp CD, Johnson JC, Cotton BA: Secondary abdominal compartment syndrome after severe extremity injury: are early, aggressive fluid resuscitation strategies to blame? *J Trauma* 2008;64:280-285

Markowitz DM, Shander A, Javidroozi M, Klein JJ, Perelman SI, Nemeth J, Ergin MS: Postoperative blood loss and transfusion associated with use of Hextend in cardiac surgery patients at a blood conservation center. *Transfusion* 2008;48:768-775

Murray CK, Hsu JR, Solomkin JS, et al: Prevention and management of infections associated with combat-related extremity injuries. *J Trauma* 2008;64:S239-S251

Murray CK: Epidemiology of infections associated with combat-related injuries in Iraq and Afghanistan. *J Trauma* 2008;64:S232-S238

Owens BD, Kragh JF, Wenke JC, Macaitis J, Wade CE, Holcomb JB: Combat wounds in Operation Iraqi Freedom and Operation Enduring Freedom. *J Trauma* 2008;64:295-299

Petersen K, Hayes DK, Blice JP, Hale RG: Prevention and management of infections associated with combat-related head and neck injuries. *J Trauma* 2008;64:S265-S276

Rhee P, Brown C, Martin M, et al: QuikClot use in trauma for hemorrhage control: case series of 103 documented uses. *J Trauma* 2008;64:1093-1099

Ritenour AE, Dorlac WC, Fang R, Woods T, Jenkins DH, Flaherty SF, Wade CE, Holcomb JB: Complications after fasciotomy revision and delayed compartment release in combat patients. *J Trauma* 2008;64:S153-S162

Ryan KL, Batchinsky A, McManus JG, Rickards CA, Convertino VA: Changes in pulse character and mental status are late responses to central hypovolemia. *Prehosp Emerg Care* 2008;12:192-198

Schwarz D, Amir L, Dichter R, Figenberg Z: The use of a powered device for intraosseous drug and fluid administration in a national EMS: a 4-year experience. *J Trauma* 2008;64:650-654

Stein DM, Dutton RP, Kramer ME, Handley C, Scalea TM: Recombinant Factor VIIa: decreasing time to intervention in coagulopathic patients with severe traumatic brain injury. *J Trauma* 2008;64:620-627

Stinger HK, Spinella PC, Perkins JG, et al: The ratio of fibrinogen to red cells transfused affects survival in casualties receiving massive transfusions at an army combat support hospital. *J Trauma* 2008;64:S79-S85

Varoga D, Drescher W, Lippross S, Pufe T, Schutz R: Nonsteroidal anti-inflammatory drug (NSAID)-related spontaneous compartment syndrome resulting from severe platelet dysfunction. *J Traum* 2008; E-pub ahead of print

Von Hoff DD, Kuhn JG, Burris HA, Miller LJ: Does intraosseous equal intravenous? A pharmacokinetic study. *Am J Emerg Med* 2008;26:31-38

Warner KJ, Copass MK, Bulger EM: Paramedic use of needle thoracostomy in the prehospital environment. *Prehosp Emerg Care* 2008;12:162-168

Wortmann GW, Valadka AB, Moores LE: Prevention and management of infections associated with combat-related central nervous system trauma. *J Trauma* 2008;64:S252-S256

Yun HC, Branstetter JG, Murray CK: Osteomyelitis in military personnel wounded in Iraq and Afghanistan. *J Trauma* 2008;64:S163-S168

## **February 2008**

Arnaud F, Tomori T, Saito R, McKeague A, Prusaczyk WK, McKarron RM: Comparative efficacy of granular and bagged formulations of the hemostatic agent QuikClot. *J Trauma* 2007;63:775-782

Borgman MA, Spinella PC, Perkins JG, Grathwohl KW, Repine T, Beekley AC, Sebesta J, Jenkind D, Wade CE, Holcomb JB: The ratio of blood products transfused in patients receiving massive transfusions at a combat support hospital. *J Trauma* 2007;63:805-813

Curtis KM, Henriques HF, Fanciullo G, Reynolds CM, Suber F: A fentanyl-based pain management protocol provides early analgesia for adult trauma patients. *J Trauma* 2007;63:819-826

Guyette FX, Wang H, Cole JS: King airway use by air medical providers. *Prehospital Emergency care* 2007;11:473-476

Harcke HT, Pearse LA, Levy AD, Getz JM, Robinson SR: Chest wall thickness in military personnel: Implications for needle thoracentesis in tension pneumothorax. *Milit Med* 2007;172:1260-1263

Kozen BG, Kircher SJ, Henao J, Godinez FS, Johnson AS: An alternative hemostatic dressing: comparison of Celox to HemCon and QuikClot. *Acad Emerg Med* 2008;15:74-81

McManus J, Hurtado T, Pusateri A, Knoop KJ: A case series describing thermal injury resulting from zeolite use for hemorrhage control in combat operations. *Prehospital Emergency Care* 2007;11:67-71

Stansbury LG, Lalliss SJ, Branstetter JG, Bagg MR, Holcomb JB: Amputations in U.S. military personnel in the current conflicts in Afghanistan and Iraq. *J Ortho Trauma* 2008;22:43-46

Zengerink I, Brink PR, Laupland KB, et al: Needle thoracostomy in the treatment of tension pneumothorax in trauma patients: What size needle? *J Trauma* 2008;64:111-114

## **November 2007**

Beekley AC, Starnes BW, Sebesta JA: Lessons Learned from modern Military Surgery. *Surg Clin N Am* 2007;87:157-184

Butler FK, Holcomb JB, Giebner SG, McSwain NE, Bagian J: Tactical Combat Casualty Care 2007: Evolving Concepts and Battlefield Experience. *Milit Med* 2007; 172(S):1-19

Colyer MH, Weber ED, Weichel ED, Dick JS, Bower KS, Ward TP, Haller JA: delayed intraocular foreign body removal without endophthalmitis during Operations Iraqi Freedom and Enduring Freedom. *Ophthalmology* 2007;114:1439-1437

Gryth D, Rocksen D, Persson JKE, et al: Severe lung contusion and death after high-velocity behind-armor blunt trauma: relation to protection level. *Milit Med* 2007;172:1110-1116

Hammett M, Watts D, Hooper T, Pearse L, Naito N: Drowning deaths of U.S. service personnel associated with motor vehicle accidents occurring in Operation Iraqi Freedom and Operation enduring freedom, 2003-2005. *Milit Med* 2007;172:875-878

Schreiber MA, Perkins J, Kiraly L, Underwood S, Wade C, Holcomb JB: Early predictors of massive transfusion in combat casualties. *J Am Coll Surg* 2007;205:541-545

Ward KR, Tiba MH, Holbert WH, Blocher CR, Draucker GT, Proffitt EK, Bowlin GL, Ivatury RR, Diegelmann RF: Comparison of a new hemostatic agent to current combat hemostatic agents in a swine model of lethal extremity arterial hemorrhage. *J Trauma* 2007;63:276-284

## **August 2007**

Eastridge BJ, Jenkins D, Flaherty S, Schiller H, Holcomb JB: Trauma system development in a theater of war: experiences from Operation Iraqi Freedom and Operation Enduring Freedom. *J Trauma* 2006;61:1366-1372

Gulam H: Care of the Muslim Patient. *Journal of Special Operations Medicine* 2005;5:39-42

Krag JF, Baer DG, Walters TJ: Extended (16-hour) tourniquet application after combat wounds; a case report and review of the literature. *J Orthop Trauma* 2007;21:274-278

Lerner EB, O'Connor RE, Schwartz R, et al: Blast-related injuries from terrorism: an international perspective. *Prehospital Emergency Care* 2007;11:137-153

Perkins JG, Schreiber MA, Wade CE, Holcomb JB: Early versus late recombinant factor VIIa in combat trauma patients requiring massive transfusion. *J Trauma* 2007;62:1099-1101

Peterson K, Riddle MS, Danko JR, Blazes DL, Hayden R, Tasker SA, Dunne JR: Trauma-related infections in battlefield casualties from Iraq. *Ann Surg* 2007;245:803-811

Rogatsky GG, Mayevsky A: The life-saving effect of hyperbaric oxygenation during early-phase severe blunt chest injuries. *Undersea Hyperb Med* 2007;34:75-81